

Original Research Article

Cardiovascular and renal oxidative stress-mediated toxicities associated with bisphenol-A exposures are mitigated by *Curcuma longa* in rats

Temitayo Olabisi Ajibade¹, Oluwaseun Olanrewaju Esan^{2,*}, Israel Mark Osawere¹, Moses Olusola Adetona³, Oluwasanmi Olayinka Aina⁴, Odunayo Ibraheem Azeez¹, Ayobami Deborah Obisesan⁵, Ademola Adetokunbo Oyagbemi¹, Olufunke Eunice Ola-Davies¹, Temidayo Olutayo Omobowale², Adebawale Benard Saba⁵, Adeolu Alex Adedapo⁵, Momoh Audu Yakubu⁶, Oluwafemi Omoniyi Oguntibeju⁷

¹Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

²Department of Veterinary Medicine, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

³Department of Anatomy, Faculty of Basic Medical Sciences, University of Ibadan, Nigeria

⁴Department of Anatomy, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

⁵Department of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

⁶Department of Environmental & Interdisciplinary Sciences, College of Science, Engineering & Technology, COPHS, Texas Southern University, Houston, TX, USA

⁷Department of Biomedical Sciences, Faculty of Health and Wellness Sciences, Cape Peninsula University of Technology, Bellville 7535, South Africa

Article history:

Received: Jun 15, 2022

Received in revised form:

Aug 16, 2022

Accepted: Aug 17, 2022

AJP, Vol. 14, No. 2, Mar-Apr 2024, 202-214.

<https://dx.doi.org/10.22038/AJP.2023.23367>

AJP.2023.23367

* Corresponding Author:

Tel: +2348077898811

esan.seun2014@gmail.com

Keywords:

Bisphenol A

Curcuma longa

Rat

Antioxidant

Anti-inflammatory

Abstract

Objective: *Curcuma longa* Rhizome (CLR), due to its potent antioxidant phytochemical constituents, was investigated for its effects on bisphenol A (BPA)-induced cardiovascular and renal damage.

Materials and Methods: Sixty rats were randomly selected, and grouped as control, BPA (100 mg/kg), BPA and CLR 100 mg/kg, BPA and CLR 200 mg/kg, CLR 100 mg/kg, and CLR 200 mg/kg for 21 days. Oxidative stress indices, antioxidant status, blood pressure parameters, genotoxicity, and immunohistochemistry were determined.

Results: Rats exposed to the toxic effects of BPA had heightened blood pressure, lowered frequency of micronucleated polychromatic erythrocytes, and decreased activities of antioxidant enzymes compared with rats treated with CLR. Moreover, administration of CLR significantly ($p < 0.05$) lowered malondialdehyde content and reduced the serum myeloperoxidase activity. Immunohistochemical evaluation revealed significantly ($p < 0.05$) increased expressions of cardiac troponin and Caspase 3 in the BPA group compared with the CLR-treated groups.

Conclusion: *C. longa* ameliorated cardiotoxic and nephrotoxic actions of bisphenol-A via mitigation of oxidative stress, hypertension, and genotoxicity.

Please cite this paper as:

Ajibade TO, Esan OO, Osawere IM, Adetona MO, Aina OO, Azeez OI, Obisesan AD, Oyagbemi AA, Ola-Davies OE, Omobowale TO, Saba AB, Adedapo AA, Yakubu MA, Oguntibeju OO. Cardiovascular and renal oxidative stress-mediated toxicities associated with bisphenol-A exposures are mitigated by *Curcuma longa* in Rats. Avicenna J Phytomed, 2024; 14(2): 202-214.

Introduction

Bisphenol A (BPA) is an inorganic environmental contaminant and endocrine disruptor found in water, air, and soil (Kang et al., 2006; Murata et al., 2018) (see Figure 1). In addition to the environmental sources of exposure, humans and animals are exposed via dietary packages containing high levels of BPA (Chen et al., 2016), with consequent toxicological manifestations on cardio-renal physiology (Gassman, 2017; Wu et al., 2021). In the cardiovascular system, BPA causes cardiac arrhythmia, myocardial infarction, and tachycardia by aggravating the formation of pro-oxidant molecules (Ola-Davies and Olukole, 2018), with consequent induction of structural and functional oxidative stress-mediated changes to deoxyribonucleic acid (Khodayar et al., 2018). Likewise, BPA promotes apoptosis, mitochondrial dysfunction, and inflammation in the renal tissue (Wu et al., 2021).




Figure 1. Schematic diagram illustrating chemical structure of bisphenol-A

Medicinal plants such as *Asparagus officinalis*, *Adiantum capillus*, Ginger, *Pistacia integerrima*, *Vincetoxicum arnottianum*, and *Murraya koenigii* have been reported to positively modulate several mammalian physiological systems and ameliorate BPA-induced toxicities in rats (Yousaf et al., 2016; Kanwal et al., 2018; Mohammed et al., 2020). Moreover, antioxidant-rich phytochemical constituents of medicinal plants such as *Asparagus officinalis* have been reported to ameliorate oxidative stress and histopathological changes in the kidney of rats intoxicated with BPA (Poormoosavi et al., 2018). In the same vein, green tea extract and epigallocatechin gallate have been reported to be effective in preventing or reducing metabolic disorders induced

by BPA due to their antioxidant and anti-inflammatory activities (Mohsenzadeh et al., 2021). *Curcuma longa* rhizome (CLR) Linn is a plant traditionally used as alternative therapy to conventional drugs for the management of diseases (Sultana et al., 2021). The biological activities of CLR for which it has been explored for the alternative management of diseases, include anti-inflammatory, anticancer, antioxidant, antibacterial, antimicrobial, antidiabetic, and immunomodulatory activities (Cárdenas-Garza et al., 2021; Mekonnen and Desta, 2021; Memarzia et al., 2021; Gururani et al., 2022; Ghasemi et al., 2022; Ghasemi et al., 2022). The antioxidant activity of curcumin, an important phytochemical constituent of CLR, reportedly ameliorated toxicant-induced organ derangements, including cardiac and renal toxicities (Ramkumar et al., 2021). The *in vivo* micronucleated polychromatic erythrocytes (MnPCEs) as a marker of genomic instability and DNA damage arising from environmental pollutants and some drugs such as BPA have been reported (Motto et al., 2020; Sommer et al., 2020; Bhilwade et al., 2014; Rocco et al., 2012). In view of the high risk of exposure to BPA from environmental sources, the present study was designed to investigate the modulatory effects of CLR against BPA-induced cardiovascular and renal toxicities. Also, we aimed to investigate the antioxidant potential of CLR against BPA-induced, oxidative stress-mediated derailment of normal physiological functioning in Wistar rats.

Materials and Methods

Experimental groupings

Sixty adult male Wistar rats having weight range 120 - 150 grams were purchased and selected without prejudice to weight or size as follows: Groups A, control, B, BPA (100 mg/kg) only, C, BPA and 100 mg/kg CLR, D, BPA and 200 mg/kg of CLR, E, 100 mg/kg of CLR only,

and group F, 200 mg/kg of CLR only. We chose the dosage of BPA from the previous study of Akintunde *et al.* (2020), while that of *C. longa* was adapted from the study of (Oridupa *et al.*, 2021). The administration of BPA alone or together with CLR was done by oral gavage, for a period 3 weeks. We followed ethical regulations in accordance with national and institutional guidelines for the protection of animal welfare during experiments.

Experimental plant

The *Curcuma longa* rhizome purchased from different Markets in Ibadan and its environs within Oyo state, Nigeria were collected freshly and preserved at optimum temperature. The CLR were identified at the department of Botany, University of Ibadan and the voucher specimen number UIH-24571 was deposited in the herbarium. The fresh rhizomes (200 g) were rinsed, air dried and then grinded into fine powder. The powder was weighed and immersed in absolute ethanol (2.5 l) for two 48 hr with regular stirring. The mixture was thereafter filtered with muslin cloth and a Whitman filter paper producing the final filtrate, which was concentrated at 45°C with a rotary evaporator, dried in a temperature-controlled oven, and stored at 4°C until required for use. The final extract weighed 72.26 g, giving a percentage yield of 12.05%.

Haemodynamic parameters

The systolic and diastolic blood pressures and mean arterial pressure were determined non-invasively in conscious animals by using an automated blood pressure monitor (CODA SI, Kent Scientific Corporation, USA) on day 20 of the experiment. The readings were taken in the quiescent state, following animal acclimatization as previously documented by Omobowale *et al.* (2019).

Serum and tissue preparation

The rats were humanely handled, and about 3 ml of blood were collected into plain bottles and subjected to centrifugation at 4,000 revolution per minute for 15 min, to obtain the serum which was carefully decanted into clean bottles, and refrigerated at -4°C throughout the experimental period.

Following anaesthesia (87.5 mg/kg Ketamine/ 12.5 mg/kg Xylazine), the heart and kidney tissues were harvested from carefully dissected rats and weighed before homogenisation in twenty (20) volumes for the heart and ten (10) volumes for the kidneys of 0.1 M phosphate buffer, and subjected to cold centrifugation at 4,000 revolution per minute for 15 min to obtain post mitochondrial fraction (Oyagbemi *et al.*, 2019).

In vivo micronucleus assay technique

The proximal ends of the femurs were carefully removed with a pair of scissors until a small opening to the marrow became visible. The femur was submerged in fetal calf serum and the marrow was flushed out gently by aspiration and flushing on glass slides. The marrow suspension was positioned on one end of a slide and spread by pulling the material behind polished cover glass held at an angle of 45°. Slides were fixed in methanol for 3-5 min; allowed to dry for 24 hr, stained with May-Gruenwald stain, and later with 5 % diluted Giemsa solution for at least thirty minutes. Slides were then rinsed in phosphate buffer for about 30 sec and in distilled water. Slides were then air-dried. The dried stained slides were then mounted in dibutylphthalate polystyrene xylene (DPX) with coverslips, and viewed under the microscope at X 100 magnification using oil immersion for the presence of micronucleated polychromatic erythrocytes (MNPCE). Scoring was done using a tally counter.

Determination of *in vivo* antioxidant status and oxidative stress indices

Markers of oxidative stress and antioxidant status such as glutathione (GSH) level were determined as described by Jollow et al. (1974). Superoxide dismutase (SOD) activity was determined as described by Misra and Fridovich (1972), with slight modification from our laboratory (Oyagbemi et al., 2019). The activity of glutathione peroxidase (GPx) was also measured according to the method of Beutler et al. (1963), while the activity of glutathione S-transferase (GST) was estimated by the method of Habig et al. (1974). Further, protein thiol (PSH) and non-protein thiol (NPSH) contents were determined as previously described by Ellman (1959).

The hydrogen peroxide (H_2O_2) production and serum myeloperoxidase (MPO) activities were estimated as reported by Wolff (1994) and Pulli et al. (2013), respectively. Vitamin C contents were quantified as previously described by Jacques-Silva et al. (2001). Malondialdehyde (MDA) content as an index of lipid peroxidation was quantified in the cardiac and renal tissues according to the method of Varshney and Kale (1990). The absorbance was estimated against blank at 532 nm. Lipid peroxidation index was calculated with a molar extinction coefficient of 1.56×10^5 M/cm. Serum nitric oxide concentrations were measured spectrophotometrically at 548 nm according to the method of Olaleye et al. (2007).

Protein determination

Tissue and serum protein concentrations were determined with Biuret method as previously described (Gornal et al., 1949). Potassium iodide was added to the reagent to prevent precipitation of Cu^{2+} ions as Cuprous oxide.

Histopathology

The heart and kidneys were collected in 10% formalin saline buffer for proper

fixation. These tissues were processed and embedded in paraffin wax. Sections of 5 – 6 μ m in thickness were made and stained with Haematoxylin and Eosin for histopathological examination (Drury et al., 1976).

Immunohistochemistry

The immunolocalization of cardiac troponin (CTnI) and caspase 3 was determined according to the method of Oyagbemi et al. (2019), with cardiac and renal tissues fixed using 10% paraformaldehyde, embedded in paraffin, and sectioned at a thickness of 5 μ m. The antigen retrieval was carried out in 10 mM citrate (6.0), in microwave oven. Goat serum (E-1R-R217A) was used to prevent specific binding. Cardiac antitroponin I (E-AB-70275: 1:500) and anticaspase 3 polyclonal antibody (E-AB-13815:1:200) were used for incubation for 2 hr, at room temperature, and thereafter, probed with secondary antibody (E-1R-R217B). Substrate diaminobenzidine (DAB) was added to improve immunopositive reactions.

Data analysis

All values were expressed as mean $\pm$ standard deviation (SD) and the test of significance between two groups was estimated with Student's t-test. The One-Way Analysis of Variance (ANOVA) with Turkey's post-hoc test was also carried out with p-values < 0.05 considered statistically significant.

Results

Effects of bisphenol-A and *C. longa* on blood pressure parameters

Bisphenol-A caused a significant ($p < 0.05$) increase in systolic, diastolic, and mean arterial pressures in the rats exposed to BPA compared with the control group. Treatment groups BPA + 100 mg/kg CLR and BPA + 200 mg/kg CLR exhibited a significantly lowered parameters of blood pressure relative to the untreated group as indicated in Figure 2.




Figure 2. Effects of BPA and CLR on blood pressure parameters; systolic, diastolic and mean arterial blood pressure. mg/kg, Bisphenol-A + CLR 200 mg/kg, CLR Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone. Values are presented as Mean $\pm$ SD (n=5).

Effects of Bisphenol A and *C. longa* on micronucleated polychromatic erythrocyte (MPCs)

As highlighted in Figure 3, Bisphenol-A caused increased frequency of MPCs in comparison with the BPA + 100 mg/kg CLR and BPA + 200 mg/kg CLR groups (Figure 3). The anti-genotoxic effect of CLR was demonstrated as indicated, with a significant reduction in the frequency of MPCs in comparison to the BPA group. It is worth noting that 200 mg/kg of CLR normalized the frequency of MPCs to near the control values (Figure 3).

Effects of *C. longa* and Bisphenol-A on organo-somatic index

BPA administration caused a significant ($p < 0.05$) reduction in body weight when compared with the control (Table 1). Similar observation was found in rats administered with BPA + 100 mg/kg CLR relative to rats administered only with BPA (Table 1). However, there was an appreciable increase in the body weights of rats administered with BPA + 200 mg/kg *C. longa*, 100 mg/kg CLR and 200 mg/kg CLR (Table 1).




Figure 3. Effects of BPA and CLR on micronucleated polychromatic erythrocyte of Wistar albino rats. Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone. Values are presented as Mean $\pm$ SD (n=5).

Cardio-renal oxidative stress indices and antioxidant status

Bisphenol-A intoxication caused a significant ($p < 0.05$) decrease in reduced glutathione (GSH) content, superoxide

dismutase (SOD), glutathione S-transferase (GST) and glutathione peroxidase (GPx) activities, in cardiac and renal tissues, compared with the control group (Tables 2 and 3). Similarly, there

Curcuma longa mitigates bisphenol-A-induced toxicity

was a noticeable improvement in GSH content of cardiac tissues which was statistically significant in rats administered with BPA +100 mg/kg CLR relative to rats administered only with BPA (Table 2). However, we also recorded a significant ($p<0.05$) decrease in GSH and vitamin C contents, and the activities of SOD, GST and GPx of rats intoxicated with 50 mg/kg

of BPA when compared to the control (Table 3). Bisphenol-A toxicity caused a significant ($p<0.05$) increase in cardiac hydrogen peroxide (H_2O_2) generation, malondialdehyde (MDA) content and significantly decreased protein thiol (PSH) and non-protein thiol (NPSH) contents, respectively, when compared to the control group.

Table 1. Effects of *Curcuma longa* and Bisphenol-A on body and organ weights in Wistar rats

Parameters	Control	Bisphenol-A (BPA) 50 mg/kg	BPA +100 mg/kg <i>Curcuma longa</i>	BPA +200 mg/kg <i>Curcuma longa</i>	100 mg/kg <i>Curcuma longa</i>	200 mg/kg <i>Curcuma longa</i>
Final Body Weight (g)	139.8±18.60	136.0±11.40 ^a	134.0±16.73 ^b	146.0±5.50 ^b	154.0±5.50 ^b	164.0±11.40 ^b
Initial Body weight (g)	98±8.36	102±13.02	104±11.40	114±8.94	112±10.95	114±8.94
Heart (g)	0.45±0.06	0.43±0.03	0.37±0.03 ^{a, b}	0.30±0.03 ^{a, b}	0.29±0.03 ^a	0.31±0.01 ^a
Kidney (g)	0.91±0.07	0.89±0.04	0.77±0.08 ^{a, b}	0.55±0.08 ^{a, b}	0.59±0.07 ^a	0.65±0.05 ^a

Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone at $p<0.05$ ($n=5$). Superscripts denote statistical difference across all groups in each row. Values are presented as mean±standard deviation.

Table 2. Effects of *Curcuma longa* and Bisphenol-A on cardiac antioxidant defense system in Wistar rats

Parameters	Control	Bisphenol-A (BPA) 50 mg/kg	BPA +100 mg/kg <i>Curcuma longa</i>	BPA +200 mg/kg <i>Curcuma longa</i>	100 mg/kg <i>Curcuma longa</i>	200 mg/kg <i>Curcuma longa</i>
GSH (μmole/mg protein)	97.10±4.26	85.17±6.02 ^a	82.60±3.29 ^a	104.4±4.50 ^{a, b}	117.19±8.97 ^a	93.43±7.85 ^a
SOD (% inhibition)	26.52±2.55	24.88±3.34	23.30±3.72	22.77±2.31 ^a	21.48±0.93 ^a	22.11±2.20 ^a
GST (CDNB-GSH complex formed/min/mg protein)	22.41±1.69	18.10±2.76 ^a	17.65±1.10 ^a	16.80±1.20 ^a	19.04±3.60 ^a	16.40±0.84 ^a
GPx (units/mg protein)	27.30±2.50	22.86±1.64 ^a	22.64±1.20 ^a	23.15±1.25 ^a	23.32±3.23 ^a	24.36±2.41 ^a
VIT C (μmol/mg protein)	13.37±1.60	12.93±1.16	13.12±1.45	11.98±1.34	12.60±0.99	12.82±1.52

Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone at $p<0.05$ ($n=5$). Superscripts denote statistical difference across all groups in each row. Values are presented as mean±standard deviation.

Table 3. Effects of *Curcuma longa* and Bisphenol-A on renal antioxidant defense system in Wistar rats

Parameters	Control	Bisphenol-A (BPA) 50 mg/kg	BPA +100 mg/kg <i>Curcuma longa</i>	BPA +200 mg/kg <i>Curcuma longa</i>	100 mg/kg <i>Curcuma longa</i>	200 mg/kg <i>Curcuma longa</i>
GSH (μmole/mg protein)	82.96±9.02	86.43±7.28	94.53±11.35	80.12±3.03	82.90±3.92	92.82±14.72
SOD (% inhibition)	35.30±3.98	12.66±1.87 ^a	38.57±4.00 ^b	30.66±1.33 ^{a, b}	39.55±7.00	28.97±4.60 ^a
GST (CDNB-GSH complex formed/min/mg protein)	23.22±4.12	12.13±2.88 ^a	32.77±6.17 ^{a, b}	27.63±5.30 ^b	36.90±5.77 ^a	35.76±3.98 ^a
GPx (units/mg protein)	36.07±5.92	40.70±5.84	46.00±6.41 ^a	43.85±4.06 ^a	48.04±5.96 ^a	53.35±5.16 ^a
VIT C (μmole/mg protein)	1.64±0.33	1.41±0.03	1.87±0.89	1.68±0.23 ^b	1.58±0.11	1.63±0.14

Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone at $p<0.05$ ($n=5$). Superscripts denote statistical difference across all groups in each row. Values are presented as mean ± standard deviation.

Interestingly, CLR co-treatment with BPA significantly lowered H₂O₂ generation, and malondialdehyde (MDA) content, with a concomitant significant increase in PSH and NPSH (Tables 4 and 5).

Effects of Bisphenol-A and *Curcuma longa* on Myeloperoxidase (MPO)

Myeloperoxidase (MPO) as a marker of inflammation was also assessed in this

study. The BPA intoxication caused a significant ($p<0.05$) increase in the MPO values of rats exposed to BPA only compared to the control and other treatment groups (Table 6). The anti-inflammatory property of CLR at 200 mg/kg showed significant reduction ($p < 0.05$) in MPO activities in comparison to BPA intoxicated group (Table 6).

Table 4. Effects of *Curcuma longa* and Bisphenol-A on markers of oxidative stress in the heart of Wistar rats

Parameters	Control	Bisphenol-A (BPA) 50 mg/kg	BPA +100 mg/kg <i>Curcuma longa</i>	BPA +200 mg/kg <i>Curcuma longa</i>	100 mg/kg <i>Curcuma longa</i>	200 mg/kg <i>Curcuma longa</i>
H ₂ O ₂ (μmole formed/min/mg protein)	15.40±0.29	16.44±0.55 ^a	12.25±0.33 ^{a, b}	15.61±1.77 ^b	13.11±0.59 ^a	13.11±2.36 ^a
PSH (μmole/mg protein)	93.02±13.23	78.60±9.25 ^a	67.03±3.05 ^{a, b}	78.43±9.70 ^a	95.31±7.93	90.95±9.24
NPSH (μmole/mg protein)	16.09±2.12	14.97±2.90	9.83±1.97 ^{a, b}	13.96±0.71	16.18±1.60	18.10±3.15
MDA (μmole formed/mg protein)	2.67±0.61	5.12±0.53 ^a	3.46±0.70 ^{a, b}	2.46±0.80 ^b	2.80±0.39	2.75±0.76

Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone at $p<0.05$ ($n=5$). Superscripts denote statistical difference across all groups in each row. Values are presented as mean±standard deviation. Abbreviations: H₂O₂ hydrogen peroxide generation (μmol/min/mg protein), MDA malondialdehyde (μmol of MDA formed/mg protein), PSH Protein thiol (μmole/mg protein), NPSH non protein thiol (μmole/mg protein)

Table 5. Effects of *Curcuma longa* and Bisphenol-A on markers of oxidative stress of the kidney of Wistar rats

Parameters	Control	Bisphenol-A (BPA) 50 mg/kg	BPA +100 mg/kg <i>Curcuma longa</i>	BPA +200 mg/kg <i>Curcuma longa</i>	100 mg/kg <i>Curcuma longa</i>	200 mg/kg <i>Curcuma longa</i>
H ₂ O ₂ (μmole formed/min/mg protein)	149.65±15.43	185.71±14.00 ^a	113.76±13.10 ^{a, b}	125.29±6.63 ^{a, b}	118.79±15.20 ^a	135.39±15.91
PSH (μmole/mg protein)	9.75±0.27	8.79±0.42 ^a	8.26±0.15 ^{a, b}	1.37±4.10 ^{a, b, c}	1.42±0.31 ^a	1.44±0.42
NPSH (μmole/mg protein)	97.48±10.6	90.85±8.48	115.83±17.13 ^{a, b}	140.56±12.43 ^{a, c}	170.10±18.65 ^a	113.7±5.34 ^{a, c}
MDA (μmole formed/mg protein)	23.82±3.96	18.41±1.77 ^a	17.33±2.83 ^a	16.55±2.16 ^a	18.60±2.56 ^a	17.91±3.48 ^a

Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone at $p<0.05$ ($n=5$). Superscripts denote statistical difference across all groups in each row. Values are presented as mean±standard deviation. Abbreviations: H₂O₂ hydrogen peroxide generation (μmol/min/mg protein), MDA malondialdehyde (μmol of MDA formed/mg protein), PSH Protein thiol (μmole/mg protein), NPSH non protein thiol (μmole/mg protein)

Table 6. Effects of *Curcuma longa* and Bisphenol-A on serum myeloperoxidase activity in Wistar rats

Parameters	Control	Bisphenol-A (BPA) 50 mg/kg	BPA +100 mg/kg <i>Curcuma longa</i>	BPA +200 mg/kg <i>Curcuma longa</i>	100 mg/kg <i>Curcuma longa</i>	200 mg/kg <i>Curcuma longa</i>
MPO (unit/mg protein)	6.90±1.40	10.91±2.16 ^a	6.12±1.01 ^b	3.63±1.54 ^{a, b}	6.68±0.81	6.46±0.25

Superscript (a) indicates a significant difference when compared to the control while (b) indicates a significant difference when compared to Bisphenol A alone at $p<0.05$ ($n=5$). Superscripts denote statistical difference across all groups in each row. Values are presented as mean±standard deviation.

Immunohistochemistry of cardiac Caspase 3

The BPA only showed the highest expression of cardiac Caspase 3 compared to all other groups. However, BPA + 100 mg/kg CLR and BPA + 200 mg/kg CLR showed a lower expression of cardiac caspase compared to the BPA only group. CLR crude extract also showed mild expression of cardiac caspase 3 (black arrows) as indicated in Figure 4.

Immunohistochemistry of cardiac Troponin I

As shown in Figure 5, BPA toxicity gave the highest expression of cardiac troponin as compared to all other groups. However, co-administration of BPA with

CLR gave a lower expression of cardiac troponin compared to the BPA only group. Also, administration of CLR alone showed mild expression of cardiac troponin (Figure 5).

Histopathological changes

In the cardiac tissues, BPA induced focal loss of myofibre striation, degeneration, and inflammatory cell infiltration. These lesions were absent in the CLR treated groups (Figure 6). In the kidneys of BPA group, mild to severe haemorrhages were observed in the renal interstitium; while rats co-administered with BPA + 100 mg/kg CLR and 200 mg/kg CLR revealed less visible pathological lesions as shown in Figure 7.




Figure 4. The immunohistochemistry of cardiac caspase 3. A (control), B (BPA), C (BPA + CLR 100 mg/kg), D (BPA + CLR 200 mg/kg), E (CLR 100 mg/kg), and F (CLR 200 mg/kg). Slides stained with high definition Haematoxylin. Magnification at X 100




Figure 5. The immunohistochemistry of cardiac troponin I. A (control), B (BPA), C (BPA + CLR 100 mg/kg), group D (BPA + CLR 200 mg/kg), Group E (CLR 100 mg/kg), and group F (CLR 200 mg/kg). Slides stained with high definition Haematoxylin. Magnification at X 100




Figure 6. The histopathology of the heart showing the effect of BPA with or without CLR in Wistar albino rats. A (control), B (BPA), C (BPA + CLR 100 mg/kg), D (BPA + CLR 200 mg/kg), E (CLR 100 mg/kg), and F (CLR 200 mg/kg). Slides stained with high definition Haematoxylin and Eosin. Magnification at X 100




Figure 7. The histopathology of the heart showing the effect of Bisphenol A with or without Curcuma longa in Wistar albino rats. A (control), B (BPA), C (BPA + CLR 100 mg/kg), D (BPA + CLR 200 mg/kg), E (CLR 100 mg/kg), and F (CLR 200 mg/kg). Slides stained with high definition Haematoxylin and Eosin. Magnification at X 100.

Discussion

The utilization of medicinal plant products for management of ill-health of humans is commonplace in the developing world due to affordability, perceived beneficial effects and reduced side effects (Laar *et al.*, 2021; Ashima *et al.*, 2021). Our results showed that CLR ameliorated hypertension associated with BPA toxicity as compared to the control. We recorded a significant reduction in systolic, diastolic, and mean arterial pressures in rats treated with CLR, indicative of anti-hypertensive action of CLR. This observation of antihypertensive effect for CLR corroborates earlier report of Hasimun *et al.* (2021) who reported CLR as a vasodilator with potent antihypertensive

effect mediated by improved vascular elasticity and potentiation of nitric oxide bioavailability. The anti-hypertensive action of CLR might be correlated with anti-inflammatory and antioxidant activity of CLR as earlier mentioned.

Also, in the present study, BPA-induced oxidative stress, seen as significant increases in the pro-oxidant but decreases in the systemic antioxidants, were mitigated by CLR. This observation underscores the widely acclaimed potent antioxidant effect of CLR which may have contributed significantly to the mitigation of BPA toxicities and hypertension seen in rats in the present study. Oxidative stress has been frequently implicated in the pathogenesis of hypertension, and its

amelioration usually correlates with normotensive status and normal vascular activities. The modulation of the vasomotor system involves reactive oxygen species as mediators of vasoconstriction induced by angiotensin II, endothelin-1 and urotensin-II, and the bioavailability of nitric oxide (NO), a major vasodilator, is highly dependent on the redox status (Rodrigo et al., 2011). Moreover, there is a positive correlation between the impairment of the antioxidant capacity and blood pressure elevation, thus underlying the relevance of antioxidant-rich molecules for the treatment of hypertension (Gonzalez et al., 2014). Therefore, mitigation of oxidative stress and normotensive status observed in the present study may be linked to the high constituents of antioxidant phytochemicals of CLR, as earlier reported (Braga et al., 2018).

From our study, we discovered that BPA intoxication significantly increased the frequency of micronucleated polychromatic erythrocytes (MnPCEs). Previous studies reported that increased frequency of MnPCEs has been positively correlated to genomic instability and damage to DNA (Motto et al., 2021; Sommer et al., 2020). Also, research findings elsewhere revealed that exposure to BPA caused DNA damage and mutagenicity (Jalal et al., 2017; Gassman et al., 2016; Pfeifer et al., 2015; Iso et al., 2006). Our finding agrees with earlier reports on the genotoxic and mutagenic effect of BPA intoxication. We found it exciting to report that rats intoxicated with BPA and treated with CLR had lower values of MnPCEs, which is indicative of antigenotoxic property of CLR in a dose-dependent manner with the 200 mg/kg of CLR being more potent. Hence, our findings are in line with previous report of antigenotoxic and non-mutagenic effects CLR (Hosseini et al., 2018; Soleimani et al., 2018; Mendonça et al., 2015).

The high immunohistochemical expression of cardiac troponin and caspase

3 indicates involvement of myocardial destruction and apoptosis in BPA-induced cardiotoxicity. High sensitive biomarkers such as cardiac troponins I (cTnI) and T (cTnT) have been shown to be highly sensitive and specific markers of myocardial cell injury (Hammarsten et al., 2018). Exposure to 50 mg/kg of BPA for 8 weeks was reported to cause morphological and structural changes in the rat myocardium in the form of vacuolated myocytes, focal loss of myofibrils, and distorted mitochondria and tubular system (Abd El-Haleem et al., 2012). Furthermore, modulation of estrogen receptor signaling, alteration of handling protein expressions, $^{+}$ cardiac Ca inhibition/activation, oxidative ion channel stress and free radical formation, and genome/transcriptome modifications have been implicated in BPA-toxicity (Bahey et al., 2019). Therefore, the attenuated immunohistochemical expressions of cardiac troponin and caspase 3, as observed in rats administered with CLR, in this study, strongly suggest anti-apoptotic mechanism of the plant in the mitigation of toxic mechanisms associated with BPA exposure. This observation corroborates an earlier report on the reversal of neuronal apoptosis by CLR in sevoflurane-induced toxicity via the p-ERK1/2 pathway (Yu et al., 2021). Moreover, mderived reactive oxygen -itochondria species were reported to be reduced in CLR extract treated SH SY5Y cells with the downregulation of sionmRNA expres associated X protein γ Bcl $\Delta\gamma$ levels of p γ and caspase (Ma et al., 2017), further underlining the positive modulatory role of CLR in the suppression of apoptotic mechanisms in toxicological processes.

In conclusion, *Curcuma longa* ameliorated the cardiotoxic and nephrotoxic effects of bisphenol A in a dose-dependent manner. Therefore, medicinal preparations containing *C. longa* could play beneficial roles in the treatment/management of toxicities

associated with environmental or dietary bisphenol A exposure.

Conflicts of interest

The authors have declared that there is no conflict of interest.

References

- Abd El-Haleem MR, Abass MA. 2012. Effect of bisphenol A on the myocardium of adult male albino rats and the possible role of lycopene: a histological and biochemical study. *Egypt J Histol*, 35: 326-339.
- Akintunde JK, Akintola TE, Adenuga GO, Odugbemi ZA, Adetoye RO, Akintunde OG. 2020. Naringin attenuates bisphenol-A mediated neurotoxicity in hypertensive rats by abrogation of cerebral nucleotide depletion, oxidative damage and neuroinflammation. *Neurotoxicology*, 81: 18-33.
- Ashima A, Jitendra G, Reena G. 2021. Miracles of herbal phytomedicines in treatment of skin disorders: natural healthcare perspective. *Infect Disord Drug Targets*, 21: 328-338.
- Bahey NG, AbdElaziz HO, Elsayed GKK. 2019. Potential toxic effect of bisphenol A on the cardiac muscle of adult rat and the possible protective effect of Omega-3: a histological and immunohistochemical study. *J Microsc Ultrastruct*, 7: 1-8.
- Beutler E, Duron O, Kelly BM. 1963. Improved method for the determination of blood glutathione. *J Lab Clin Med*, 61: 882-888.
- Bhilwade HN, Jayakumar S, Chaubey RC. 2014. Age-dependent changes in spontaneous frequency of micronucleated erythrocytes in bone marrow and DNA damage in peripheral blood of Swiss mice. *Mutat Res Genet Toxicol Environ Mutagen*, 770: 80-84.
- Braga MC, Vieira ECS, de Oliveira TF. 2018. *Curcuma longa* L. leaves: characterization (bioactive and antinutritional compounds) for use in human food in Brazil. *Food Chem*, 265: 308-315.
- Cárdenas-Garza GR, Elizondo LJH, Bazaldúa RAF, Chávez MA, Pérez HRA, Martínez DAJ, López VSM, Rodríguez RJ, Sánchez CRM, Castillo VU, Rodríguez LOE. 2021. Benefits of cardamom (*Elettaria cardamomum* (L.) Maton) and turmeric (*Curcuma longa* L.) extracts for their applications as natural anti-inflammatory adjuvants. *Plants*, 10: 1908.
- Drury RA, Wallington EA. 1976. Carlton's histopathological techniques, pp. 139-142, London, Oxford University Press.
- Ellman GL. 1959. Tissue sulfhydryl groups. *Arch Biochem Biophys*, 82: 70-77.
- Gassman NR, Coskun E, Jaruga P, Dizdaroglu M, Wilson SH. 2016. Combined effects of high-dose Bisphenol A and oxidizing agent (kbro3) on cellular microenvironment, gene expression, and chromatin structure of Ku70-deficient mouse embryonic fibroblasts. *Environ Health Perspect*, 24: 1241-1252.
- Ghasemi SZ, Memarzia A, Behrouz S, Gholamnezhad Z, Boskabady MH. 2022. Comparative effects of *Curcuma longa* and curcumin on paraquat-induced systemic and lung oxidative stress and inflammation in rats. *Avicenna J Phytomed*, 12: 414-424.
- Gornal AG, Bardawill JC, David MM. 1949. Determination of serum proteins by means of biuret reaction. *Res J Biol Chem*, 177: 751-766.
- Gururani S, Gairola K, Kumar R, Prakash O, Dubey SK. 2022. Altitudinal and geographical variations in phytochemical composition and biological activities of *Curcuma longa* accession from Uttarakhand, the Himalayan region. *J Food Proc Preserv*, 46: e16384.
- Habig WH, Pabst MJ, Jakoby WB. 1974. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J Biol Chem*, 249: 7130-7139.
- Hammarsten O, Mair J, Möckel M, Lindahl B, Jaffe AS. 2018. Possible mechanisms behind cardiac troponin elevations. *Biomarkers*, 23: 725-734.
- Hasimun P, Sulaeman AH, Yani M. 2021. Effect of *Curcuma longa* L. extract on noninvasive cardiovascular biomarkers in hypertension animal models. *J Appl Pharmaceut Sci*, 11: 085-089.
- Hosseini A, Hosseinzadeh H. 2018. Antidotal or protective effects of *Curcuma longa* (turmeric) and its active ingredient, curcumin, against natural and chemical toxicities: A review. *Biomed Pharmacother*, 99: 411-421.
- Iso T, Watanabe T, Iwamoto T, Shimamoto A, Furuichi Y. 2006. DNA damage caused by

- bisphenol A and estradiol through estrogenic activity. *Biol Pharm Bull*, 29: 206-210.
- Jacques-Silva MC, Nogueira CW, LC, Flores, Rocha JBT. 2001. Diphenyl diselenide and ascorbic acid changes deposition of selenium and ascorbic acid in liver and brain of mice. *Pharmacol Toxicol*, 88: 119-125.
- Jalal N, Surendranath AR, Pathak JL, Yu S, Chung CY. 2017. Bisphenol A (BPA) the mighty and the mutagenic. *Toxicol Rep*, 16: 76-84.
- Jollow DJ, Mitchel JR, Zampaglione N. 1974. Bromobenzene-induced liver necrosis protective role of glutathione and evidence for 3, 4-bromobenzene oxide as the hepatotoxic metabolite. *Pharmacology*, 11: 151-169.
- Kang JH, Katayama Y, Kondo F. 2006. Biodegradation or metabolism of bisphenol A: from microorganisms to mammals. *Toxicology*, 217: 81-90.
- Khodayar MJ, Kalantari H, Mahdavinia M, Khorsandi L, Alboghobeish S, Samimi A, Alizadeh S, Zeidooni L. 2020. Protective effect of naringin against BPA-induced cardiotoxicity through prevention of oxidative stress in male Wistar rats. *Drug Chem Toxicol*, 43: 85-89.
- Laar A, Ampah EA, Fernandez Y, Amevinya GS, Nortey P, Benyah F, Akamah J, Ambenne M, Lamptey P, Free C, Legido-Quigley H, Perel P. 2021. What the herbal medicine can do for me in a week, the orthodox does in a year: Perceived efficacy of local alternative therapies influences medication adherence in patients with atherosclerotic cardiovascular disease. *Health Expect*, 24: 444-455.
- Ma XW, Guo RY. 2017. Dose-dependent effect of *Curcuma longa* for the treatment of Parkinson's disease. *Exp Ther Med*, 13: 1799-1805.
- Mekonnen A, Desta W. 2021. Comparative study of the antioxidant and antibacterial activities of *Rumex abyssinicus* with commercially available *Zingiber officinale* and *Curcuma longa* in Bahir Dar city, Ethiopia. *Chem Biol Technol Agric*, 8: 2.
- Memarzia A, Khazdair MR, Behrouz S, Gholamnezhad Z, Jafarnejhad M, Saadat S, Boskabady MH. 2021. Experimental and clinical reports on anti-inflammatory, antioxidant, and immunomodulatory effects of *Curcuma longa* and curcumin, an updated and comprehensive review. *Biofactors*, 47: 311-350.
- Mendonça LM, Machado Cda S, Teixeira CC, Freitas LA, Bianchi ML, Antunes LM. 2015. Comparative study of curcumin and curcumin formulated in a solid dispersion: Evaluation of their antigenotoxic effects. *Genet Mol Biol*, 38: 490-498.
- Misra HP, Fridovich I. 1972. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem*, 247: 3170-3175.
- Mohsenzadeh MS, Razavi BM, Imenshahidi M, Tabatabaee Yazdi SA, Mohajeri SA, Hosseinzadeh H. 2021. Potential role of green tea extract and epigallocatechin gallate in preventing bisphenol A-induced metabolic disorders in rats: Biochemical and molecular evidence. *Phytomedicine*, 92: 153754.
- Motto EA, Lawson-Evi P, Kantati Y, Eklug-Gadegbeku K, Aklikokou K, Gbeassor M. 2020. Antihyperglycemic activity of total extract and fractions of *Anogeissus leiocarpus*. *J Drug Deliv Therapeut*, 10: 107-113.
- Ola-Davies OE, Olukole SG. 2018. Gallic acid protects against bisphenol A-induced alterations in the cardio-renal system of Wistar rats through the antioxidant defense mechanism. *Biomed Pharmacother*, 107: 1786-1794.
- Olaleye SB, Adaramoye OA, Erigbali PP, Adeniyi OS. 2007. Lead exposure increases oxidative stress in the gastric mucosa of HCl/ethanol exposed rats. *World J Gastroenterol*, 13: 5121-5126.
- Oridupa OA, Oyagbemi AA, Adejumo O, Falade FB, Obisesan AD, Abegunde BA, Ekwem PC, Adegboye VO, Omobowale TO. 2021. Compensatory depression of arterial pressure and reversal of ECG abnormalities by *Annona muricata* and *Curcuma longa* in hypertensive Wistar rats. *J Complement Integr Med*, 19: 375-382.
- Oyagbemi AA, Omobowale TO, Awoyomi OV, Ajibade TO, Falayi OO, Ogunpolu BS, Okotie UJ, Asenuga ER, Adejumo OA, Hassan FO, Ola-Davies OE, Saba AB, Adedapo AA, Yakubu MA. 2019. Cobalt

- chloride toxicity elicited hypertension and cardiac complication via induction of oxidative stress and upregulation of COX-2/Bax signaling pathway. *Hum Exp Toxicol*, 38: 519-532.
- Pfeifer D, Chung YM, Hu MC. 2015. Effects of low-dose bisphenol A on DNA damage and proliferation of breast cells: The role of c-Myc. *Environ Health Perspect*, 23: 1271-1279.
- Poormoosavi MS, Najafzadehvarzi H, Behmanesh MA, Amirgholam R. 2018. Protective effects of *Asparagus officinalis* extract against Bisphenol A-induced toxicity in Wistar rats. *Toxicol Reports*, 5: 427-433.
- Ramkumar S, Thulasiram HV, RaviKumar A. 2021. Improvement in serum amylase and glucose levels in diabetic rats on oral administration of bisdemethoxycurcumin from *Curcuma longa* and limonoids from *Azadirachta indica*. *J Food Biochem*, 45: 13674.
- Rocco L, Peluso C, Stingo V. 2012. Micronucleus test and comet assay for the evaluation of zebrafish genomic damage induced by erythromycin and lincomycin. *Environ Toxicol*, 27: 598-604.
- González J, Valls N, Brito R, Rodrigo R. 2014. Essential hypertension and oxidative stress: New insights. *World J Cardiol*, 6: 353-366.
- Rodrigo R, González J, Paoletto F. 2011. The role of oxidative stress in the pathophysiology of hypertension. *Hypertens Res*, 34: 431-440.
- Soleimani V, Sahebkar A, Hosseinzadeh H. 2018. Turmeric (*Curcuma longa*) and its major constituent (curcumin) as nontoxic and safe substances: Review. *Phytother Res*, 32: 985-995.
- Sommer S, Buraczewska I, Kruszewski M. 2020. Micronucleus Assay: The state of art, and future directions. *Int J Mol Sci*, 21: 1534.
- Sultana S, Munir N, Mahmood Z, Riaz M, Akram M, Rebezov M, Kuderinova N, Moldabayeva Z, AliShariati M, Rauf A, Rengasamy RR. 2021. Molecular targets for the management of cancer using *Curcuma longa* Linn. phytoconstituents: A review. *Biomed Pharmacother*, 135: 111078.
- Varshney R, Kale RK. 1990. Effect of calmodulin antagonists on radiation induced lipid peroxidation in microsomes. *Int J Radiat Biol*, 58: 733-743.
- Wolff SF. 1994. Ferrous ion oxidation in the presence of ferric ion indicator xylenol orange for measurement of hydrogen peroxides. *Methods Enzymol*, 233: 182-189.
- Wu B, Zhao Q, Li Z, Min Z, Shi M, Nie X, He Q, Gui R. 2021. Environmental level bisphenol A accelerates alterations of the reno-cardiac axis by the MAPK cascades in male diabetic rats: An analysis based on transcriptomic profiling and bioinformatics. *Environ Pollut*, 287: 117671.
- Pulli B, Ali M, Forghani R, Schob S, Hsieh KL, Wojtkiewicz G, Linnoila JJ, Chen JW. 2013. Measuring myeloperoxidase activity in biological samples. *PLoS One*, 8: 67976.
- Yousaf B, Amina LG, Wang R, Qadir A, Ali MU, Kanwal Q, Munir B, Asmatullah, Abbas Z. 2016. Bisphenol-A exposure and healing effects of *Adiantum capillus-veneris* L. plant extract (APE) in bisphenol A-induced reproductive toxicity in albino rats. *Environ Sci Pollut Res Int*, 23: 11645-11657.
- Yu Z, Yan Y, Lou Y. 2021. *Curcuma longa* Linn extract suppresses neuronal apoptosis induction by sevoflurane via activation of the ERK1/2 pathway. *Tropical J Pharmaceut Res*, 20: 269-274.