


Ayesha Jacobs* and Sosthene Nyomba Kamanda

Crystal structure of tris(piperazine-1,4-dium) bis(2-hydroxy-1,2,3-propane-tricarboxylate) pentahydrate, $C_{24}H_{56}N_6O_{19}$



<https://doi.org/10.1515/ncrs-2025-0330>

Received July 31, 2025; accepted September 4, 2025;
published online September 19, 2025

Abstract

$C_{24}H_{56}N_6O_{19}$, triclinic, $P\bar{1}$ (no. 2), $a = 6.0132(12)$ Å, $b = 11.667(2)$ Å, $c = 13.152(3)$ Å, $\alpha = 67.10(3)^\circ$, $\beta = 88.54(3)^\circ$, $\gamma = 83.18(3)^\circ$, $V = 843.8(3)$ Å³, $Z = 2$, $R_{gt}(F) = 0.0606$, $wR_{ref}(F^2) = 0.1336$, $T = 173(2)$ K.

CCDC no.: 2484891

The molecular structure is shown in the figure. Table 1 contains the crystallographic data and the list of the atoms including atomic coordinates and displacement parameters can be found in the cif-file attached to this article.

1 Source of materials

The 2-hydroxy-1,2,3-propane-tricarboxylic acid (citric acid) monohydrate was obtained from UniLAB Saarchem (Merck) and piperazine hexahydrate from BDH laboratories. Diethyl ether was obtained from Merck. The citric acid monohydrate was dried in an oven at 60 °C for 2 h. Citric acid (0.26 mmol, 0.050 g) and piperazine hexahydrate (0.26 mmol, 0.052 g) was dissolved in 2.0 mL of a 1:1 mixture of diethyl

*Corresponding author: Ayesha Jacobs, Faculty of Applied Sciences, Chemistry Department, Cape Peninsula University of Technology, PO Box 1906, Bellville 7535, South Africa, E-mail: jacobsa@cput.ac.za. <https://orcid.org/0000-0002-5508-3291>

Sosthene Nyomba Kamanda, Faculty of Applied Sciences, Chemistry Department, Cape Peninsula University of Technology, PO Box 1906, Bellville 7535, South Africa

Table 1: Data collection and handling.

Crystal:	Colourless block
Size:	0.27 × 0.17 × 0.11 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.12 mm ⁻¹
Diffractometer, scan mode:	Bruker Apex-II diffractometer, φ and ω scans
$\theta_{\max}$, completeness:	28.3°, 100 %
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$, R_{int} :	13092, 4187, 0.026
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 3,830
$N(\text{param})_{\text{refined}}$:	278
Programs:	Bruker, ^{1,2} WinGX, ³ SHELX, ^{4,5} X-Seed ⁶

ether:water with slight heating. The solution was allowed to evaporate at room temperature. Block crystals were obtained after 5 months of slow evaporation.

2 Experimental details

The hydroxyl hydrogen atom, water hydrogen atoms and amine hydrogens were located in the difference electron density map and freely refined with isotropic temperature factors. The C–H atoms were geometrically constrained at 0.99 Å with $U_{\text{iso}}(\text{H}) = 1.2 \text{ Å}^2$.

3 Comment

Citric acid is naturally found in citrus fruits, however commercially it is sourced from fermentation processes primarily using the fungus *Aspergillus niger*.^{7,8} Citric acid has uses in a wide variety of industries including food, pharmaceutical and chemical industries.^{9,10} Recently, cocrystallisation studies of citric acid with selected heterocyclic compounds have been reported.¹¹ Piperazine is an anthelmintic drug for veterinary use.¹² Piperazine has been used in cocrystallisation studies involving natural products and active pharmaceutical ingredients.^{13–15}

The crystal structure of piperazine citrate monohydrate, (C₄H₁₂N₂)²⁺(C₆H₆O₇)²⁻·H₂O in the monoclinic space group *Pn* ($a = 9.2055(12) \text{ Å}$, $b = 6.8314(9) \text{ Å}$, $c = 11.2443(14) \text{ Å}$, $\beta = 112.047(2)^\circ$, $V = 655.41(15) \text{ Å}^3$) has been reported by Liu.¹⁶ The asymmetric unit of (C₄H₁₂N₂)²⁺(C₆H₆O₇)²⁻·H₂O contains a double deprotonated citrate anion, a doubly protonated piperazine and one water molecule.

The asymmetric unit of the current structure, 3(C₄H₁₂N₂)²⁺·2(C₆H₅O₇)³⁻·5(H₂O), contains a triple deprotonated citrate anion, 1.5 piperazinium cations and 2.5 water molecules. The piperazinium cations lie on centres of inversion and

all three of the carboxylic acids of citric acid are deprotonated. There are several NH $\cdots$ O, OH $\cdots$ O and CH $\cdots$ O hydrogen bonds in the structure. The piperazinium cations and water molecules are hydrogen bonded to the carboxylate groups of the citrate anion. The hydroxyl group of the citrate forms an intramolecular OH $\cdots$ O contact with one of the carboxylate oxygens. There are numerous hydrogen bonded rings and chains. In particular, two citrate and two piperazinium ions form R_4^4 (24) rings involving four NH $\cdots$ O heterosynthons. Citrate and piperazinium ions are linked via NH $\cdots$ O hydrogen bonds creating C_4^4 (26) chains. The two and a half water molecules and one citrate ion produce R_4^4 (10) rings via OH $\cdots$ O hydrogen bonds. The water molecules occupy channels parallel to [100] with different water environments present including donor $\cdots$ donor (DD), donor $\cdots$ donor $\cdots$ acceptor (DDA) and donor $\cdots$ donor $\cdots$ acceptor $\cdots$ acceptor (DDAA).

Author contributions: All the authors have accepted responsibility for the entire content of this submitted manuscript and approved submission.

Conflict of interest: The authors declare no conflicts of interest regarding this article.

References

1. Bruker. APEX2, SAINT-PLUS and SADABS; Bruker AXS Inc.: Madison, Wisconsin, USA, 2009.
2. Bruker. XPREP; Bruker AXS Inc.: Madison, Wisconsin, USA, 2004.
3. Farrugia, L. J. WinGX and ORTEP for Windows: An Update. *J. Appl. Cryst.* **2012**, 45, 849–854.
4. Sheldrick, G. M. A Short History of SHELX. *Acta Cryst.* **2008**, A64, 112–122.
5. Sheldrick, G. M. Crystal Structure Refinement with SHELXL. *Acta Cryst.* **2015**, C71, 3–8.
6. Barbour, L. J. X-Seed-A Software Tool for Supramolecular Crystallography. *Supramol. Chem.* **2001**, 1, 189–191.
7. Kubicek, C. P.; Röhr, M.; Rehm, H. J. Citric Acid Fermentation. *Crit. Rev. Biotechnol.* **1985**, 3, 331–373.
8. Angumeenal, A. R.; Venkappayya, C. D. An Overview of Citric Acid Production. *LWT-Food Sci. Technol.* **2013**, 50, 367–370.
9. Ciriminna, R.; Meneguzzo, F.; Delisi, R.; Pagliaro, M. Citric Acid: Emerging Applications of Key Biotechnology Industrial Product. *Chem. Cent. J.* **2017**, 11, 22.
10. Lambros, M.; Tran, T.; Fei, Q.; Nicolaou, M. Citric Acid: A Multifunctional Pharmaceutical Excipient. *Pharmaceutics* **2022**, 14, 972.
11. Easmin, S.; Pedireddi, V. R. Systematic Exploration of Structural Topologies in Hydrogen-Bonded Supramolecular Assemblies of Citric Acid with Different Heterocyclic Compounds. *ACS Omega* **2023**, 8, 23202–23217.
12. Liu, M.; Panda, S. K.; Luyten, W. Plant-Based Natural Products for the Discovery and Development of Novel Anthelmintics Against Nematodes. *Biomolecules* **2020**, 10, 426.
13. Ungur, D. T.; Iordache, C. A.; Brăilă, C. A.; Pop, D. A.; David, M. C.; Pandele-Cuşu, J.; Fruth, V.; Pongratz, I.; Iacob, B.-C.; Bodoki, E.; Pop, M. M. Structural Insights into the Resveratrol-Piperazine Cocrystal

- Forms Enabling the Cocrystallization Process Development from Solution. *Cryst. Growth Des.* **2025**, 25, 1330–1343.
14. Wang, Z.; Li, S.; Li, Q.; Wang, W.; Liu, M.; Yang, S.; Zhang, L.; Yang, D.; Du, G.; Lu, Y. A Novel Cocrystal of Daidzein with Piperazine to Optimize the Solubility, Permeability and Bioavailability of Daidzein. *Molecules* **2024**, 29, 1710.
15. Andre, V.; Braga, D.; Grepioni, F.; Duarte, M. T. Crystal Forms of the Antibiotic 4-aminosalicylic Acid: Solvates and Molecular Salts with Dioxane, Morpholine, and Piperazine. *Cryst. Growth Des.* **2009**, 9, 5108–5116.
16. Liu, L.-L. Piperazine-1,4-dium 2-(carboxymethyl)-2-hydroxybutanedioate Monohydrate. *Acta Cryst.* **2010**, E66, o2191.