

RESEARCH ARTICLE

Clofibrate, a PPAR- α agonist, abrogates sodium fluoride-induced neuroinflammation, oxidative stress, and motor incoordination via modulation of GFAP/Iba-1/anti-calbindin signaling pathways

Ademola A. Oyagbemi¹  | Olamide E. Adebisi¹ | Kabirat O. Adigun¹ |
 Blessing S. Ogunpolu² | Olufunke O. Falayi³ | Fasilat O. Hassan¹ |
 Oluwabusayo R. Folarin^{4,5} | Adedeji K. Adebayo³ | Olumuyiwa A. Adejumo² |
 Ebunoluwa R. Asenuga^{4,6} | Olufunke E. Ola-Davies¹ | Temidayo O. Omobowale² |
 James O. Olopade^{5,6} | Adebowale B. Saba³ | Adeolu A. Adedapo³ |
 Sanah M. Nkadimeng⁷ | Lyndy J. McGaw⁷ | Oluwafemi O. Oguntibeju⁸ |
 Momoh A. Yakubu⁹

¹Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

²Department of Veterinary Medicine, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

³Department of Veterinary Pharmacology and Toxicology, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

⁴Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Benin, Nigeria

⁵Department of Medical Laboratory Science, College of Health Sciences, Ladoké Akintola University of Technology, Osogbo, Osun State, Nigeria

⁶Department of Veterinary Anatomy, Faculty of Veterinary Medicine, University of Ibadan, Nigeria

⁷Phytomedicine Programme, Department of Paraclinical Sciences, University of Pretoria, Faculty of Veterinary Science, Onderstepoort, South Africa

⁸Phytomedicine and Phytochemistry Group, Department of Biomedical Sciences, Faculty of Health and Wellness Sciences, Cape Peninsula University of Technology, Bellville, South Africa

⁹Department of Environmental & Interdisciplinary Sciences, College of Science, Engineering & Technology, Texas Southern University, Houston, Texas

Correspondence

Dr. Ademola A. Oyagbemi, Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Ibadan, Nigeria.
 Email: ademola.oyagbemi778@gmail.com

Funding information

National Research Foundation; Cape Peninsula University of Technology

Abstract

Fluoride is an environmental contaminant that is ubiquitously present in air, water, and soil. It is commonly added in minute quantity to drinking water, toothpaste, and mouth rinses to prevent tooth decay. Epidemiological findings have demonstrated that exposure to fluoride induced neurodevelopmental toxicity, developmental neurotoxicity, and motor disorders. The neuroprotective effect of clofibrate, a peroxisome proliferator-activated receptor alpha agonist, was investigated in the present study. Forty male Wistar rats were used for this study and randomly grouped into 10 rats per group as control, sodium fluoride (NaF) alone (300 ppm), NaF plus clofibrate (250 mg/kg), and NaF plus lisinopril (10 mg/kg), respectively, for 7 days. NaF was administered in drinking water while clofibrate and lisinopril were administered by oral gavage. Markers of neuronal inflammation and oxidative stress, acetylcholinesterase activity, and neurobehavioral (hanging wire and open field) tests were performed. Immunohistochemistry was performed on brain tissues, and they were probed with glial fibrillary acidic protein, ionized calcium-binding adaptor molecule 1,

and cerebellar Ca^{2+} -binding protein calbindin-D28k. The results showed that NaF significantly increased oxidative stress and neuroinflammation and inhibited AChE activity. Immunostaining showed reactive astrocytes, microgliosis, loss of dendritic spines, and arborization in Purkinje cells in rats administered only NaF. Neurobehavioral results showed that cotreatment of NaF with clofibrate improved muscular strength and locomotion, reduced anxiety, and significantly reduced astrocytic count. Overall, cotreatment of NaF with either clofibrate or lisinopril showed neuroprotective effects by mitigating neuronal inflammation and oxidative and motor incoordination. Hence, clofibrate could be seen as a novel drug candidate against neurodegeneration and motor disorders.

KEYWORDS

clofibrate, neuroinflammation, neuroprotection, oxidative stress, sodium fluoride

1 | INTRODUCTION

Fluoride is an environmental contaminant that is ubiquitously present in air, water, and soil.¹ Meanwhile, oxidative stress and free radical generation have been reported to play a fundamental role in the pathogenesis of sodium fluoride (NaF) toxicity.^{2–4} However, few studies have documented the toxic effects of NaF on the central nervous system (CNS) and motor incoordination.^{5–9}

Recently, there has been a significant increase in the global incidence of hypertension and neurobehavioral deficit such as motor incoordination.¹⁰ Hypertension is a known important risk factor for the development of Alzheimer disease (AD) and cerebral small vessel disease.¹¹ Hypertension, on the other hand, has been documented to enhance deposition of beta-amyloid ($\text{A}\beta$) protein, which can ultimately trigger AD and aging-related dementia.¹¹ In other findings, attention-deficit hyperactivity disorder has been reported in spontaneously hypertensive rats.^{12–14}

The involvement of oxidative stress and free radical generation has been linked to the pathophysiology of NaF-induced neurodegeneration.¹⁵ Previous research reports have documented neuronal apoptosis, developmental neurotoxicity in rat offspring, cytoskeleton damage and decreased signal, excessive endoplasmic reticulum stress and autophagic dysfunction, synaptic impairment, and developmental neurotoxicity following exposure to NaF.^{5,7,8,16,17} Furthermore, fragmentation and redistribution of mitochondria away from the axons of the cortical neurons¹⁸ decreased the expressions of glucose transporter 1 (GLUT1) and glial fibrillary acidic protein (GFAP),¹⁹ induction of S-phase cell-cycle arrest, upregulation of NF-kappa B and DNA damage in primary rat hippocampal neurons,²⁰ behavioral deficit,⁹ alteration of cognition, emotion, and synaptic plasticity in rat hippocampus.^{21,22}

Clofibrate is a peroxisome proliferator-activated receptor alpha (PPAR- α) agonist.²³ The novel neuroprotective effect of PPAR- α agonists such as pioglitazone and fenofibrate against neuroinflammation, neuronal damage, and axonal injury has been reported via antioxidant and anti-inflammatory properties.^{24–26} However, there is a paucity of information on the novel neuroprotective properties of clofibrate;

hence, this study seeks to unravel the neuroprotective effects of clofibrate and its molecular mechanism of action against NaF-induced neurotoxicity and neurobehavioral deficit.

2 | MATERIALS AND METHODS

2.1 | Chemicals

Clofibrate, lisinopril, trichloroacetic acid, sodium hydroxide, O-dianisidine, hydrogen peroxide (H_2O_2), xylene orange (XO), potassium hydroxide, reduced glutathione (GSH), oxidized glutathione, NaF, thiobarbituric acid, and 1,2-dichloro-4-nitrobenzene were purchased from Sigma (St. Louis, Missouri). Normal goat serum, biotinylated antibody, and horse radish peroxidase system were purchased from KPL Inc (Gaithersburg, Maryland), anti-glial fibrillary acidic protein (GFAP; 1:1000) rabbit polyclonal antibodies from Dako (Denmark), anti-calbindin (1:12000) rabbit monoclonal antibody D-28k from Swant (Switzerland), and anti-ionized calcium-binding adapter molecule-1 (Iba-1; 1:800) rabbit monoclonal antibody from Wako, while 3,3'-diaminobenzidine (DAB, Wako, Richmond, USA) tablets were purchased from Amresco LLC (Ohio). All other chemicals used for this study were of analytical grade.

2.1.1 | Experimental animals and design

In the study, forty male Wistar rats were randomly grouped into 10 rats per group as control, NaF alone (300 ppm), NaF plus clofibrate (250 mg/kg), and NaF plus lisinopril (10 mg/kg), respectively, for 7 days. NaF was administered using drinking water while clofibrate and lisinopril were administered by oral gavage. The animals were also fed with rat cubes ad libitum. The rats were kept in wire mesh cages under controlled light cycle (12 hours light/12 hours dark) and fed with commercial rat chow ad libitum and water was supplied liberally. The blood of the rats was taken on the eighth day and rats were euthanized on the ninth day. All the animals received humane care according to the criteria outlined in the *Guide for the Care and Use of Laboratory Animals*.

2.2 | Hanging wire test

To test for the effect of treatments of rats on muscular strength, a hanging wire test was used. At the end of the treatment, rats were placed individually with their forepaws on a horizontally suspended wire (1 mm in diameter) and placed 47 cm above a soft foam landing area. Both ends of the hanging wire were tightly secured. Each of the experimental animals was placed with its forepaws on the wire. Rats were timed from the moment they were placed on the wire until they dropped from the wire. This reflected muscular strength in rats.²⁷ The hanging wire was cleaned between each rat using 70% alcohol. Each rat was tested three times.

2.3 | Open field test

The open field apparatus was a white box measuring 72 × 72 × 36 cm. One of the sides was covered with Plexiglass so that rats could be visible in the apparatus. Black lines divided the floor into sixteen 18 × 18 cm squares with a central square (18 × 18 cm) at the middle of the open field.²⁸ Rats were placed at the center of the open field and allowed to explore the box for 5 minutes. Line crossing, center square entries, center square duration, rearing, stretched attend posture (SAP), grooming, freezing, and fecal pellet were recorded as previously described.²⁹ The open field maze (OFM) was cleaned between each rat using 70% alcohol.

2.4 | Blood sample collection and plasma preparation

Approximately 3 mL of blood was collected by retro-orbital venous puncture using heparinized capillary tubes into plain bottles and allowed to clot. The clotted blood was then centrifuged at 4000 rpm for 10 minutes. Clear serum was separated using a Pasteur pipette into another plain tube and then stored at 4°C until required.

2.5 | Preparation of neuronal homogenates

At the end of treatment, rats were euthanized and the whole brain was excised, rinsed and homogenized using 50 mM Tris-HCl buffer (pH 7.4) containing 1.15% KCl. The homogenates were subjected to cold centrifugation at 4°C at a speed of 10 000 g for 15 minutes. The post mitochondrial fractions (PMFs) obtained from brain homogenates were used for biochemical assays.

2.6 | Biochemical assays

2.6.1 | Neuronal makers of oxidative stress

H₂O₂ generation was determined according to the method of Wolff.³⁰ The reaction mixture was subsequently incubated at room temperature for 30 minutes. The mixtures were read at an absorbance of 560 nm and H₂O₂ generated was extrapolated from the H₂O₂ standard curve. The malondialdehyde (MDA) content as an index of lipid peroxidation was quantified according to the method of Varshney and Kale.³¹ The absorbance was measured against a blank of distilled water at 532 nm. Lipid peroxidation was calculated with a molar extinction coefficient of

1.56 × 10⁵ M⁻¹ cm⁻¹. The absorbance of the sample was measured at 370 nm. Vitamin C contents were measured as earlier described.³² The protein and non-protein thiol (NPSH) contents were determined as described by Ellman.³³ The activity of acetylcholinesterase (AChE) was measured as described by Ellman.³³

2.6.2 | Neuronal antioxidant defense system

The superoxide dismutase (SOD) assay was carried out using the method of Misra and Fridovich with slight modifications from our laboratory.^{34,35} The increase in absorbance at 480 nm was monitored every 30 seconds for 150 seconds. The one unit of SOD activity was given as the amount of SOD necessary to cause 50% inhibition of the oxidation of adrenaline to adrenochrome. GSH was estimated by the method of Jollow et al.³⁶ Glutathione peroxidase (GPx) activity was also measured according to Beutler et al.³⁷ Glutathione S-transferase (GST) was estimated by the method of Habig et al.³⁸ using 1-chloro-2,4-dinitrobenzene as the substrate. The decrease in absorbance at 340 nm was measured, and enzymatic activity was calculated with a molar extinction coefficient of 6.1 mmol/L⁻¹ cm. Protein carbonyl (PCO) contents in the brain tissues were measured using the method of Reznick and Packer.³⁹ Protein concentration was determined using the Biuret method of Gornal et al.⁴⁰ using bovine serum albumin as a standard.

2.7 | Immunohistochemical staining

The immunolocalization of cerebrum and cerebellum was described as earlier reported¹ to determine the expressions of GFAP for astrocytes (cerebral cortex), ionized calcium-binding adaptor molecule 1 (Iba1) for microglia (cerebral cortex), and cerebellar Ca²⁺-binding protein calbindin-D28k (CB) in the brain (cerebellum). Sections were observed using a light microscope (Leica LAS-EZ) with Leica software application suite version 3.4 equipped with a digital camera. Immunoreactivity was quantified using the Image J software by quantifying the total tissue area and the specific area of immunolocalization or immunopositivity. This was then followed by dividing the immunopositive reactions with total tissue area and then expressed as arbitrary units.

2.8 | Statistical analysis

Data obtained were analyzed with one-way analysis of variance followed by Tukey's test to compare the mean of one group with the other. All values are expressed as mean ± SD. The test of significance between two groups was estimated by the Student's *t* test.

3 | RESULTS

3.1 | Behavioral studies

3.1.1 | Hanging wire test

The hanging wire test was used to evaluate the motor function disorder induced by NaF. The hanging latency values in the control group were 99.92 ± 12.09 seconds, NaF alone of 26.39 ± 4.93 seconds, NaF

plus clofibrate (250 mg/kg) of 82.39 ± 4.93 seconds, and NaF plus lisinopril (10 mg/kg) of 80.56 ± 12.13 seconds. The rats treated with NaF alone had significantly ($P \leq .05$) reduced hanging latency compared with other groups (Figure 1). The hanging latency of the NaF plus clofibrate and NaF plus lisinopril groups was similar to that of the control group. Rats coadministered with clofibrate or lisinopril and NaF held onto the wire for a longer duration while 90% of the rats administered with NaF alone fell before mid-way into the test.

3.1.2 | Open field maze

Neither the neurotoxicant nor the test compounds modified ambulatory activity compared with the control group in the center square, line crossing, freezing, and hanging wire (Figure 1). Similar results were observed in center square duration and rearing (Figure 1). NaF alone or in combination with clofibrate significantly increased the SAP when compared with the control and cotreatment with lisinopril (Figure 2). Furthermore, in contrast to administration of NaF alone, every other group showed a significant decrease in rearing frequency in the OFM (Figure 2). The effects produced by lisinopril in the OFM were similar to those observed in the control group.

3.2 | AChE activity and neuronal antioxidant defense system

The results as indicated in figures showed that NaF significantly inhibited AChE activity in comparison with the control and rats cotreated with clofibrate or lisinopril (Figure 2). The content of neuronal GSH fell significantly following the administration of NaF while

a significant increase in GSH content was observed in rats coadministered with either clofibrate or lisinopril (Figure 3). The MDA content as an index of lipid peroxidation was found to increase significantly in the rats that received only NaF relative to the control (Figure 3). However, on the other hand, clofibrate or lisinopril coadministration led to a significant decrease in the values of MDA as shown in Figure 3. Similarly, the neuronal H_2O_2 generation and PCO content increased significantly in NaF alone treated rats as compared to control and other treatment groups (Figure 3). Interestingly, there was reduction in the activity of brain GPx. However, the reduction was not statistically significant across groups (Figure 4). The GST activity in the animals that received only NaF reduced significantly while improvement in the GST activity was recorded in the group that received NaF in combination with lisinopril (Figure 4). The activity of SOD increased significantly in groups that received NaF in combination with either clofibrate or lisinopril (Figure 4). The administration of NaF caused a significant increase in neuronal activity to control nitric oxide (NO) bioavailability while coadministration with either clofibrate or lisinopril caused a significant reduction in neuronal NO content (Figure 5). The protein thiol (PSH) content increased significantly in rats administered NaF when compared to the control, which was significantly reduced in groups that received either clofibrate or lisinopril along with NaF (Figure 5). Furthermore, the NPSH content was not significantly affected by the treatment as indicated in Figure 5. However, there was a significant increase in vitamin C levels in rats that received clofibrate along with NaF while NaF alone or lisinopril in combination with NaF did not alter the vitamin C content (Figure 5).




FIGURE 1 The effect of NaF on neurobehavioral tests. Alphabet "a" indicates significant difference when compared to the control at $P < .05$ while "b" indicates significant difference when compared to NaF only group at $P < .05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 2 The effect of NaF on neurobehavioral tests and acetylcholinesterase (AChE) activity. Alphabet "a" indicates significant difference when compared to the control at $P < .05$ while "b" indicates significant difference when compared to NaF only group at $P < .05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 3 The effect of NaF on markers of oxidative. Alphabet "a" indicates significant difference when compared to the control at $P < .05$ while "b" indicates significant difference when compared to NaF only group at $P < 0.05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]

3.3 | Immunohistochemistry

The immunohistochemistry of anti-calbindin showed a reduction in size of Purkinje cells in NaF alone treated rats relative to the control or groups that received either clofibrate or lisinopril in combination

with NaF (Figure 6). However, the architecture of the Purkinje cells was slightly preserved in groups that received NaF in combination with either clofibrate or lisinopril (Figure 6). On the other hand, GFAP expressions showed significant astrocytosis in NaF alone treated rats




FIGURE 4 The effect of sodium fluoride (NaF) on the antioxidant defense system. Alphabet "a" indicates significant difference when compared to the control at $P < .05$ while "b" indicates significant difference when compared to NaF only group at $P < 0.05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]




FIGURE 5 The effect of sodium fluoride (NaF) on non-enzymatic antioxidant and nitric oxide content. Alphabet "a" indicates significant difference when compared to the control at $P < .05$ while "b" indicates significant difference when compared to NaF only group at $P < 0.05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]

relative to the control or groups that received either clofibrate or lisinopril combination with NaF (Figure 7). However, there was a significant reduction in the number of astrocytes in groups that received

NaF in combination with either clofibrate or lisinopril (Figure 8). The immunolocalization of Iba1 showed significant microgliosis in NaF alone treated rats relative to the control (Figure 8). The cotreatment





FIGURE 6 The immunolocalization of anti-calbindin on sodium fluoride (NaF)-induced neurotoxicity in the cerebellum. NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride. Alphabet "a" indicates significant difference when compared to the control while "b" indicates significant difference when compared to NaF only group at $P < .05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]



with either clofibrate or lisinopril caused a significant reduction in microgliosis (Figure 8).

4 | DISCUSSION

The inhibition of AChE has been reported as one of the cardinal points of neurotoxicity in both aquatic and mammalian toxicity.⁴¹⁻⁴⁴ AChE inhibition can lead to autonomic nervous crisis resulting in both sympathetic and parasympathetic symptoms.⁴⁵ Neuropsychiatric disorders have been reported following exposure to organophosphates; these are major inhibitors of AChE.^{46,47} AD is an advanced neurodegenerative disorder that is associated with AChE activation and amyloid plaque deposition.^{48,49}

From our results, NaF significantly inhibited AChE activity; this might be the hallmark of neurotoxicity observed in this study. Furthermore, coadministration with either clofibrate or lisinopril reactivates AChE inhibition to near normal activity. AChE reactivators have been reported as antidotes and treatment of choice for organophosphorus organophosphate pesticide intoxication.^{46,47,50} Hence, clofibrate or lisinopril might be a novel cholinesterase reactivator in the near future.

Increasing evidence has shown the involvement of oxidative stress in the pathogenesis of Parkinson disease (PD) and potential drugs can be developed through modulation and inhibition of AChE activity or its antagonist.⁵¹⁻⁵⁴ The results obtained from the present study demonstrated the enhancement of oxidative stress by NaF intoxication as indicated by significant depletion of neuronal GSH, vitamin C, NPSH, SOD, and GST activities together with significant enhancement of neuronal MDA, H_2O_2 generated, and PSH and NO contents. The exaggerated production of lipid peroxidation products and depletion of enzymatic and non-enzymatic antioxidant defense system observed in this study contributed significantly oxidative stress, significantly to oxidative stress, neuroinflammation, and neurodegeneration. Our study is in tandem with previous studies that reported on how oxidative stress and neuroinflammation contributed to neurodegeneration and motor incoordination.⁵⁵ The novel antioxidant activity of clofibrate and lisinopril was demonstrated by a significant improvement in antioxidant defense system and mitigation of oxidative stress as indicated by significant reduction in markers of oxidative stress. The development of neurodegenerative disorders such as AD and PD has been positively correlated to oxidative stress and neuroinflammation.^{56,57} Similarly, oxidative fatty acid degradation in the

FIGURE 7 The immunolocalization of glial fibrillary acidic protein (GFAP) on sodium fluoride (NaF)-induced neurotoxicity in the cerebral cortex. Alphabet "a" indicates significant difference when compared to the control while "b" indicates significant difference when compared to NaF only group at $P < .05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]







FIGURE 8 The immunohistochemistry of ionized calcium-binding adapter molecule 1 (IBA1) on sodium fluoride (NaF)-induced neurotoxicity in the cerebral cortex. Alphabet "a" indicates significant difference when compared to the control while "b" indicates significant difference when compared to NaF only group at $P < .05$ ($n = 10$). NaF (300 ppm), clofibrate (250 mg/kg), and lisinopril (10 mg/kg). Ctrl, control; NaF, sodium fluoride [Color figure can be viewed at wileyonlinelibrary.com]

brain has been reported as the signaling pathway in the development of Alzheimer-like pathology.⁵⁶ Therefore, enhanced production of neuronal NO in the present study played a major role in the observable neuroinflammation. Interestingly, coadministration of clofibrate or lisinopril with NaF significantly attenuated excessive production of NO, which was indicative of anti-inflammatory properties of clofibrate and lisinopril. Lisinopril is a classic inhibitor of angiotensin-converting enzyme (ACE).

OFM is commonly used to measure mobility and anxiety-like behavior in animal models.⁵⁸ The different treatment protocols did not have any effect on the ambulatory ability of the rats; however, the animals moved freely in the box as displayed by similarity in the number of lines crossed across over the treatment groups. Rearing, an exploratory behavior, used as a measure of anxiety in the OFM increased in the NaF group, suggesting an increased anxiety level.⁵⁹ This anxiogenic potential of NaF is further confirmed by the significant increase in SAP. This indicates the reluctance of the rats to move reflecting an approach-avoidance tendency.⁶⁰ On the contrary, lisinopril administration significantly decreased this tendency showing its anxiogenic potential. The anxiety-stimulating effect of NaF reported in this study is in consonance with reports by Yuksel.⁶¹ who in addition documented an increased number of defecations in the OFM following administration.

The immunohistochemistry of anti-calbindin revealed a significant loss of Purkinje fibers following NaF intoxication. Anti-calbindin is a 28KDa protein located in the cerebellar cortex.⁶² Previous studies have demonstrated link between Purkinje cell loss and neurobehavioral derangement such as motor incoordination and essential brain tremor of which PD is a sequela.⁶³⁻⁶⁶ There is a growing evidence that Purkinje cell loss is associated with essential tremor.⁶⁷ The contribution of NaF to Purkinje cell loss in the present study was restored to near normal following coadministration of either clofibrate or lisinopril, which was indicative of their neuroprotective effects, thereby maintaining cerebellar architectural integrity. This study therefore supports a previous report on the possible neuroprotective of ACE inhibitor through the downregulation of kinin B1 receptors in the peripheral nervous system, which is responsible for neuropathic pain.⁶⁸ Oxidative stress and neuroinflammation might have contributed to the observable distortion and significant reduction in Purkinje cells in rats exposed to NaF alone.

The main intermediate filament protein in mature astrocytes is GFAP and is the critical component of the cytoskeleton during development.⁶⁹ Astrocytes are abundant in the brain and are the first reactive cells after any injury; thus, GFAP has been used as biomarker in several neurotoxicity studies.^{70,71} The significant increase in intermediate filament expressions, cellular hypertrophy, and proliferation observed in the cerebral cortex administered with NaF alone is indicative of astrocyte response to CNS insults. The astrocytic reaction is suggestive of defense against the observed increased oxidative stress induced by NaF. Astrocytes contain high concentrations of antioxidants upregulating glutathione following oxidative stress in a bid to restore CNS homeostasis.⁷²⁻⁷⁴ On the other hand, astrocytosis with hypertrophied processes is compensatory for neuronal and synaptic

degeneration as typified by the observed Purkinje cell degeneration in the rats.⁷⁵ Administration of either clofibrate or lisinopril was able to reduce astrocytic activation and expressions of GFAP, confirming their anti-inflammatory and neuroprotective properties as peroxisome proliferator-activated receptor- γ (PPAR γ) agonist.⁷⁶

In the present study, there was a significant increase in the expressions of Iba1. The Iba1 is a novel calcium-binding protein and is specifically expressed in microglia, and it has been reported to play a significant role in regulation of the function of microglia. Microglia constitute approximately 5% of all glial cells in the CNS. Activation of microglia commonly occurs in the early response of the CNS to a wide variety of pathological stimuli, including trauma, inflammation, degeneration, and ischemia.⁷⁷ They help in the elimination of microorganisms and deleterious debris by producing neurotrophic factors that are involved in neuroprotection.⁷⁸⁻⁸⁰ On the other hand, microglia also exert a cytotoxic function by releasing reactive oxygen species, NO, or inflammatory cytokines, which ultimately triggered neuronal damage.⁸¹ From the present study, administration of NaF led to a significant activation of microglia, and this might be associated with exaggeration exaggerated release of neuronal NO in NaF treated rats as compared to rats coadministered with either clofibrate or lisinopril. Therefore, clofibrate and lisinopril demonstrated their anti-inflammatory properties by mitigating microgliosis as observed in NaF alone. In conclusion, administration of NaF precipitated oxidative stress, Purkinje neuron degeneration, motor incoordination, and depletion of antioxidant defense system, while coadministration with either clofibrate or lisinopril attenuated oxidative stress, improved antioxidant defense status, and offered neuroprotection. Combining all, our findings demonstrate the neuroprotective effects of clofibrate and lisinopril as indicated in their ability to prevent astrocytosis, microgliosis, and Purkinje cell loss.

ACKNOWLEDGMENTS

We acknowledge the Neuroscience laboratory for their technical assistance and provision of antibodies. The authors are grateful to Cape Peninsula University of Technology and National Research Foundation (South Africa) for financial support granted to Prof OO Oguntibeju.

CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

ORCID

Ademola A. Oyagbemi  <https://orcid.org/0000-0002-8996-8610>

REFERENCES

1. Oyagbemi AA, Omobowale TO, Asenuga ER, et al. NaF induces hypertension and cardiac complications through generation of

- reactive oxygen species and activation of nuclear factor kappa beta. *Environ Toxicol*. 2017;32:1089-1101.
2. Oyagbemi AA, Omobowale TO, Ola-Davies EO, et al. Quercetin attenuates hypertension induced by NaF via reduction in oxidative stress and modulation of HSP 70/ERK/PPAR γ signaling pathways. *Biofactors*. 2018;44:465-479.
 3. Oyagbemi AA, Omobowale TO, Ola-Davies EO, et al. Luteolin-mediated Kim-1/NF- κ B/Nrf2 signaling pathways protects NaF-induced hypertension and cardiovascular complications. *Biofactors*. 2018;44:518-531.
 4. Oyagbemi AA, Omobowale TO, Ola-Davies EO, et al. Ameliorative effect of Rutin on NaF-induced hypertension through modulation of Kim-1/NF- κ B/Nrf2 signaling pathway in rats. *Environ Toxicol*. 2018;33:1284-1297.
 5. Chen J, Niu Q, Xia T, et al. ERK1/2-mediated disruption of BDNF-TrkB signaling causes synaptic impairment contributing to fluoride-induced developmental neurotoxicity. *Toxicology*. 2018;410:222-230.
 6. Tu W, Zhang Q, Liu Y, et al. Fluoride induces apoptosis via inhibiting SIRT1 activity to activate mitochondrial p53 pathway in human neuroblastoma SH-SY5Y cells. *Toxicol Appl Pharmacol*. 2018;347:60-69.
 7. Niu Q, Chen J, Xia T, et al. Excessive ER stress and the resulting autophagic flux dysfunction contribute to fluoride-induced neurotoxicity. *Environ Pollut*. 2018;233:889-899.
 8. Chen L, Ning H, Yin Z, et al. The effects of fluoride on neuronal function occurs via cytoskeleton damage and decreased signal transmission. *Chemosphere*. 2017;185:589-594.
 9. Adedara IA, Abolaji OA, Idris UF, et al. Neuroprotective influence of taurine on fluoride-induced biochemical and behavioral deficits in rats. *Chem Biol Interact*. 2017;261:1-10.
 10. Jassim AH, Inman DM. Evidence of hypoxic glial cells in a model of ocular hypertension. *Invest Ophthalmol Vis Sci*. 2019;60:1-15.
 11. Jeon SY, Byun MS, Yi D, Lee JY, et al. KBASE research group. Influence of hypertension on brain amyloid deposition and Alzheimer's disease signature neurodegeneration. *Neurobiol Aging*. 2019;75:62-70.
 12. Cho HS, Baek DJ, Baek SS. Effect of exercise on hyperactivity, impulsivity and dopamine D2 receptor expression in the substantia nigra and striatum of spontaneously hypertensive rats. *J Exerc Nutr Biochem*. 2014;18(4):379-384.
 13. Ko IG, Kim SE, Kim TW, et al. Swimming exercise alleviates the symptoms of attention-deficit hyperactivity disorder in spontaneous hypertensive rats. *Mol Med Rep*. 2013;8(2):393-400.
 14. Kim H, Heo H, Kim DH, et al. Treadmill exercise and methylphenidate ameliorate symptoms of attention deficit/hyperactivity disorder through enhancing dopamine synthesis and brain-derived neurotrophic factor expression in spontaneous hypertensive rats. *Neurosci Lett*. 2011;504(1):35-39.
 15. França CF, Vianna LM. Effectiveness of B vitamins on the control of hypertension and stroke events of SHRSP rats. *J Diet Suppl*. 2010;7(1):71-77.
 16. Wu J, Cheng M, Liu Q, et al. Protective role of tert-Butylhydroquinone against sodium fluoride-induced oxidative stress and apoptosis in PC12 cells. *Cell Mol Neurobiol*. 2015;35(7):1017-1025.
 17. Zhu YP, Xi SH, Li MY, et al. Fluoride and arsenic exposure affects spatial memory and activates the ERK/CREB signaling pathway in offspring rats. *Neurotoxicology*. 2017;59:56-64.
 18. Lou DD, Guan ZZ, Liu YJ, et al. The influence of chronic fluorosis on mitochondrial dynamics morphology and distribution in cortical neurons of the rat brain. *Arch Toxicol*. 2013;87(3):449-457.
 19. Jiang C, Zhang S, Liu H, et al. Low glucose utilization and neurodegenerative changes caused by NaF exposure in rat's developmental brain. *Neuromolecular Med*. 2014;16(1):94-105.
 20. Zhang M, Wang A, Xia T, He P. Effects of fluoride on DNA damage, S-phase cell-cycle arrest and the expression of NF- κ B in primary cultured rat hippocampal neurons. *Toxicol Lett*. 2008;179(1):1-5.
 21. Liu F, Ma J, Zhang H, et al. Fluoride exposure during development affects both cognition and emotion in mice. *Physiol. Behav*. 2014;124:1-7.
 22. Qian W, Miao K, Li T, Zhang Z. Effect of selenium on fluoride-induced changes in synaptic plasticity in rat hippocampus. *Biol Trace Elem Res*. 2013;155(2):253-260.
 23. Pan M, Jiao S, Reinach PS, et al. Opposing effects of PPAR α Agonism and antagonism on refractive development and form deprivation myopia in Guinea pigs. *Invest Ophthalmol Vis Sci*. 2018;59(15):5803-5815.
 24. Moreno S, Cerù MP. In search for novel strategies towards neuroprotection and neuroregeneration: is PPAR α a promising therapeutic target? *Neural Regen Res*. 2015;10(9):1409-1412.
 25. Gray E, Ginty M, Kemp K, Scolding N, Wilkins A. Peroxisome proliferator-activated receptor- α agonists protect cortical neurons from inflammatory mediators and improve peroxisomal function. *Eur J Neurosci*. 2011;33(8):1421-1432.
 26. Gray E, Ginty M, Kemp K, Scolding N, Wilkins A. The PPAR- γ agonist pioglitazone protects cortical neurons from inflammatory mediators via improvement in peroxisomal function. *J Neuroinflammation*. 2012;9:63.
 27. Soazo M, García G. Vanadium exposure through lactation produces ehavioural alterations and CNS myelin deficit in neonatal rats. *Neurotoxicol Teratol*. 2007;29(4):503-510.
 28. Carrey N, McFadyen MP, Brown RE. Effects of subchronic methylphenidate hydrochloride administration on the locomotor and exploratory behavior of prepubertal mice. *J Child Adolesc Psychopharmacol*. 2000;10(4):277-286.
 29. Brown RE, Corey SC, Moore AK. Differences in measures of exploration and fear in MHC-congenic C57BL/6J and B6-H-2K mice. *Behav Genetics*. 1999;26:263-271.
 30. Wolff SF. Ferrous ion oxidation in the presence of ferric ion indicator xylenol orange for measurement of hydrogen peroxides. *Methods Enzymol*. 1994;233:182-189.
 31. Varshney R, Kale KK. Effect of calmodulin antagonists on radiation induced lipid peroxidation in microsomes. *Int J Radiat Biol*. 1990;58(5):733-743.
 32. Jacques-Silva MC, Nogueira CW, Broch LC, Flores EMM, RochaNote JBT. Diphenyl diselenide and ascorbic acid changes deposition of selenium and ascorbic acid in liver and brain of mice. *Pharmacol Toxicol*. 2001;88(3):119-125.
 33. Ellman GL. Tissue sulfhydryl groups. *Arc Biochem Biophys*. 1959;82:70-77.
 34. Misra HP, Fridovich I. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *J Biol Chem*. 1972;247(10):3170-3175.
 35. Oyagbemi AA, Omobowale TO, Akinrinde AS, et al. Lack of reversal of oxidative damage in renal tissues of lead acetate-treated rats. *Environ Toxicol*. 2015;30(11):1235-1243.
 36. Jollow DJ, Mitchell RM. N, Bromobenzene-induced liver necrosis. Protective role of glutathione and evidence for 3, 4-bromobenzene oxide as the hepatotoxic metabolite. *Pharmacol*. 1974;11:151-169.
 37. Beutler E, Duron O, Kelly BM. Improved method for the determination of blood glutathione. *J Lab Clin Med*. 1963;61:882-888.
 38. Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases. The first enzymatic step in mercapturic acid formation. *J Bio Chem*. 1974;249(22):7130-7139.
 39. Reznick AZ, Packer L. Oxidative damage to proteins: spectrophotometric method for carbonyl assay. *Methods Enzymol*. 1994;233:357-363.

40. Gornal AG, Bardawill JC, David MM. Determination of serum proteins by means of biuret reaction. *J Biol Chem.* 1949;177(2): 751-766.
41. Serafini S, de Freitas SC, Baldissera MD, et al. Fish exposed to water contaminated with eprinomectin show inhibition of the activities of AChE and Na⁺/K⁺-ATPase in the brain, and changes in natural behavior. *Chemosphere.* 2019;223:124-130.
42. Hinojosa HG, Gutiérrez-Praena D, Prieto AI, et al. Neurotoxicity induced by microcystins and cylindrospermopsin: a review. *Sci Total Environ.* 2019;668:547-565.
43. Hauser-Davis RA, Lopes RP, Zioli RL. Inhibition of mullet (*M. liza*) brain acetylcholinesterase activity by in vitro polycyclic aromatic hydrocarbon exposure. *Mar Pollut Bull.* 2019;140:30-34.
44. Leris L, Kalogianni E, Tsangaris C, et al. Acute and sub-chronic toxicity bioassays of olive mill wastewater on the eastern mosquito fish *Gambusia holbrooki*. *Ecotoxicol Environ Saf.* 2019;175:48-57.
45. Lataro RM, Silva MAB, Mestriner FL, Cau SBA, Tostes RCA, Salgado HC. Chronic treatment with acetylcholinesterase inhibitors attenuate vascular dysfunction in spontaneously hypertensive rats. *Am J Hypertens.* 2019;32(6):579-587.
46. Jokanović M. Neurotoxic effects of organophosphorus pesticides and possible association with neurodegenerative diseases in man: A review. *Toxicol.* 2018;410:125-131.
47. Faria NM, Fassa AG, Meucci RD, et al. Occupational exposure to pesticides, nicotine and minor psychiatric disorders among tobacco farmers in southern Brazil. *Neurotoxicology.* 2014;45:347-354.
48. Saleh H, Sadeghi L. Investigation of THDOC effects on pathophysiological signs of Alzheimer's disease as an endogenous neurosteroid: inhibition of acetylcholinesterase and plaque deposition. *Bratisl Lek Listy.* 2019;120(2):148-154.
49. Hsieh YC, Guo C, Yalamanchili HK, et al. Tau-Mediated Disruption of the Spliceosome Triggers Cryptic RNA Splicing and Neurodegeneration in Alzheimer's Disease. *Cell Rep.* 2019;29(2):301-316.
50. Kovarik Z, Maček Hrvat N, Kalisiak J, et al. Counteracting tabun inhibition by reactivation by pyridinium aldioximes that interact with active center gorge mutants of acetylcholinesterase. *Toxicol Appl Pharmacol.* 2019;372:40-46.
51. Mohammadzadeh L, Hosseinzadeh H, Abnous K, Razavi BM. Neuroprotective potential of crocin against malathion-induced motor deficit and neurochemical alterations in rats. *Environ Sci Pollut Res Int.* 2018; 25(5):4904-4914.
52. Fadly EDA, Elzahha PA, Tramarin A, et al. Tackling neuroinflammation and cholinergic deficit in Alzheimer's disease: multi-target inhibitors of cholinesterases, cyclooxygenase-2 and 15-lipoxygenase. *Eur J Med Chem.* 2019;167:161-186.
53. Pan H, Zhang J, Wang Y, et al. Linarin improves the dyskinesia recovery in Alzheimer's disease zebrafish by inhibiting the acetylcholinesterase activity. *Life Sci.* 2019;222:112-116.
54. Hepnarova V, Korabecny J, Matouskova L, et al. The concept of hybrid molecules of tacrine and benzyl quinolone carboxylic acid (BQCA) as multifunctional agents for Alzheimer's disease. *Eur J Med Chem.* 2018;150:292-306.
55. Chen Y, Wu T, Li H, et al. DI-3-n-butylphthalide exerts dopaminergic neuroprotection through inhibition of Neuroinflammation. *Front Aging Neurosci.* 2019;11:44.
56. Furman R, Axelsen PH. The effects of omega-3 fatty acid deficiency during development on oxidative fatty acid degradation during maturity in a mouse model of Alzheimer's disease. *Neurobiol Aging.* 2019; 79:66-74.
57. Schmidt HL, Garcia A, Izquierdo I, Mello-Carpes PB, Carpes FB. Strength training and running elicit different neuroprotective outcomes in a β -amyloid peptide-mediated Alzheimer's disease model. *Physiol Behav.* 2019;206:206-212.
58. Prut L, Belzung C. The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: a review. *Eur J Pharm.* 2003;463:3-33.
59. Ennaceur A. Tests of unconditional anxiety – pitfalls and disappointments. *J Phys Behav.* 2014;135:55-71.
60. Molewijk HE, van der Poel AM, Olivier B. The ambivalent behaviour "stretched approach posture" in the rat as a paradigm to characterize anxiolytic drugs. *Psychopharmacol.* 1995;121:81-90.
61. Yuksel K. Effects of fluoride on anxiety and depression in mice. *Fluoride.* 2012;45:302-306.
62. Aguiar J, Fernández J, Aguilar A, et al. Ubiquitous expression of human SCA2 gene under the regulation of the SCA2 self-promoter cause specific Purkinje cell degeneration in transgenic mice. *Neurosci Lett.* 2006;392(3):202-206.
63. Louis ED. Linking essential tremor to the cerebellum: neuropathological evidence. *Cerebellum.* 2016;15(3):235-242.
64. Babij R, Lee M, Cortés E, Vonsattel JP, Faust PL, Louis ED. Purkinje cell axonal anatomy: quantifying morphometric changes in essential tremor versus control brains. *Brain.* 2013;136(10): 3051-3061.
65. Shill HA, Adler CH, Beach TG. Pathology in essential tremor. *Parkinsonism Relat Disord.* 2012;18(1):S135-S137.
66. Rajput HA, Robinson CA, Rajput ML, Robinson SL, Rajput A. Essential tremor is not dependent upon cerebellar Purkinje cell loss. *Parkinsonism Relat Disord.* 2012;18(5):626-628.
67. Labiano-Fontcuberta A, Benito-León J. Essential tremor and Parkinson's disease: are they associated? *Rev Neurol.* 2012;55(8): 479-489.
68. Kaur P, Muthuraman A, Kaur M. The implications of angiotensin-converting enzymes and their modulators in neurodegenerative disorders: current and future perspectives. *ACS Chem Neurosci.* 2015;6(4): 508-521.
69. Middeldorp J, Hol EM. GFAP in health and disease. *Prog Neurobiol.* 2011;93(3):421-443.
70. Lei J, Gao G, Feng J, et al. Glial fibrillary acidic protein as a biomarker in severe traumatic brain injury patients: a prospective cohort study. *Crit Care.* 2015;19:362.
71. McMahon PJ, Panczykowski DM, Yue JK, et al. TRACK-TBI investigators. Measurement of the glial fibrillary acidic protein and its breakdown products GFAP-BDP biomarker for the detection of traumatic brain injury compared to computed tomography and magnetic resonance imaging. *J Neurotrauma.* 2015;32(8): 527-533.
72. Dringen R. Metabolism and functions of glutathione in brain. *Prog Neurobiol.* 2000;62:649-671.
73. Chen Y, Chan PH, Swanson RA. Astrocytes overexpressing Cu, Zn superoxide dismutase has increased resistance to oxidative injury. *Glia.* 2000;33:343-347.
74. Liberto CM, Albrecht PJ, Herx LM, Yong VW, Levison SW. Pro-regenerative properties of cytokine-activated astrocytes. *J Neurochem.* 2004;89:1092-1100.
75. Gyls KH, Fein JA, Yang F, Wiley DJ, Miller CA, Cole GM. Synaptic changes in Alzheimer's disease: increased amyloid-beta and gliosis in surviving terminals is accompanied by decreased PSD-95 fluorescence. *Am J Pathol.* 2004;165:1809-1817.
76. Lee EY, Lee JE, Park JH, Shin IC, Koh HC. Rosiglitazone, a PPAR- γ agonist, protects against striatal dopaminergic neurodegeneration induced by 6-OHDA lesions in the substantia nigra of rats. *Toxicol Lett.* 2012;213(3):332-344.
77. Collins JM, King AE, Woodhouse A, Kirkcaldie MTK, Vickers JC. The effect of focal brain injury on beta-amyloid plaque deposition, inflammation and synapses in the APP/PS1 mouse model of Alzheimer's disease. *Exp Neurol.* 2015;267:219-229.

78. Kittl M, Jakab M, Steininger ST, et al. A swelling-activated chloride current in microglial cells is suppressed by Epac and facilitated by PKA - impact on phagocytosis. *Cell Physiol Biochem*. 2019;52(5):51-969.
79. Bogie JF, Stinissen P, Hendriks JJ. Macrophage subsets and microglia in multiple sclerosis. *Acta Neuropathol*. 2014;128(2):191-213.
80. Zindler E, Zipp F. Neuronal injury in chronic CNS inflammation. *Best Pract Res Clin Anaesthesiol*. 2010;24(4):551-562.
81. Chu F, Shi M, Zheng C, et al. The roles of macrophages and microglia in multiple sclerosis and experimental autoimmune encephalomyelitis. *J Neuroimmunol*. 2018;318:1-7.

How to cite this article: Oyagbemi AA, Adebisi OE, Adigun KO, et al. Clofibrate, a PPAR- α agonist, abrogates sodium fluoride-induced neuroinflammation, oxidative stress, and motor incoordination via modulation of GFAP/Iba-1/anti-calbindin signaling pathways. *Environmental Toxicology*. 2020; 35:242–253. <https://doi.org/10.1002/tox.22861>