

International Conference Proceedings

**33rd JOHANNESBURG International Conference on “Chemical,
Biological and Environmental Engineering” (JCBEE-22)**

**March 17-18, 2022
Johannesburg (South Africa)**

Editors:

Prof. Dr. S. K. Ntwampe,
Prof. Dr. E. Fosso-Kankeu
Prof. Dr. B. Topcuoğlu

ISBN – 978-989-53228-9-3

Published by

Pilares D'Elegância, Lda

Organized by



Growth and Yield of *Brassica oleracea* in Response to Phosphorus Fertilizer Application

N Mfeka¹, M Tanga¹, A.T Adetunji¹ and F. B Lewu¹,

Abstract— Cabbage is an important vegetable that is grown around the world because of its nutritional content and for medicinal purposes. Plant nutrients can deplete due to continuous cultivation, this necessitates the addition of fertilizer to sustain production. Therefore, the objective of this experiment was to investigate the effect of phosphorus fertilizer on the vegetative and reproductive growth of cabbage growth in the Western Cape. A two-season experiment was conducted using a complete randomized complete block design (RCBD), with four replicates. Treatments of phosphorus (Triple superphosphate) at four levels 109, 163.5, 218, and 222.5kg/ha. Data collection on the vegetative and yield was recorded and analyzed using the SPSS software. The results showed that treatment did not have a significant difference ($p>0.05$) on vegetative and yield and generally p fertilizer at 218 kg/ha increased all the measured parameters. An increase in yield during the last season of the experiment was also noted.

Keywords— *Brassica oleracea*, plant nutrients, phosphorus

I. INTRODUCTION

Cabbage (*Brassica oleracea*, L.), is one of the most common vegetables grown in South Africa (SA) and its production is generally concentrated in Mpumalanga and Kwazulu-Natal [2]. In SA this leafy vegetable is typically cultivated in smallholder plots and provides dietary diversity and nutrients in poor rural communities [7]–[8]. *Brassica oleracea* is rich in minerals and vitamins such as A, B1, B2, C [14], and these characteristics are an indication of good quality [15]. According to Nikzad [9], cabbage contains phosphorus which assists in the utilization of calcium and the assimilation of carbohydrates and fats in the human body. Cabbage is adaptable to a wide range of climatic conditions, soil type and nutritional benefits [2], [5]–[6]. This vegetable crop is regarded as a heavy feeder as it requires large quantities of the macronutrients [2], [4]–[5]. Natural occurring

nutrients easily get depleted over the years of cultivation this necessitates the addition of synthetic fertilizer such as phosphorus. This macro-nutrient is essential for plant growth, metabolic processes and is required plants in large quantities [5]. However, majority of research done has mainly focused on macronutrient nitrogen and its effect on the cabbage vegetative parameters and yield [4], [11]–[13]. The objective of this experiment was, therefore, to investigate the effect of phosphorus fertilizer on vegetative and reproductive growth of cabbage growth in the Western Cape.

II. MATERIALS AND METHODS

A. Study Area

The research was conducted at a Research and Teaching Farm, the Agricultural-Hub, Department of Agriculture, Wellington Campus, Cape Peninsula University of Technology, Western Cape Province. The area is located in the northern part of Wellington with the following coordinate ($S33^{\circ} 37' E19^{\circ} 37'$), with the Mediterranean climate (with hot dry summers and wet cold winters). The area receives an annual rainfall of 585 mm and approximately one-third falls between March and August. Prior to transplanting the field was tilled and plots were prepared accordingly. After transplanting uniform irrigation was supplied and the supply of irrigation water was uniform throughout the growing season. Cultivation practices such as weeding was done manually when necessary. Nitrogen and potassium fertilizer was applied and kept constant. The initial soil results indicated that the field was a loam soil with a pH of 4.6, P 23 and K 178 mg/kg and the second year analysis pH was 4.6, P 32 and K 93.9 mg/kg [10].

B. Experimental Layout and treatment application

The trial was conducted during the summer season 2019-2020. The experimental design for each year was laid in a complete randomized complete block design (RCBD), with four replicates. Each unit plot measured 3.5m x 1m, 1m between plots and plant spacing of 60cm. The treatments consisted of four rates of phosphorus (Triple superphosphate 20%): 109(T1), 163.5 (T2), 218 (T3) and 222.5 (T4) kg/ha.

C. Data collection

Data collection on plant height, number of leaves and leaf width, leaf length were recorded over a period of two weeks interval during the growth period of 2019 and 2020, data

N Mfeka is with the ¹Department of Agriculture, Faculty of Applied Science, Cape Peninsula University of Technology, P.O. Box XB Wellington 7654, South Africa (corresponding author's phone: 0218645505; email: mfekan@cput.ac.za)

M Tanga is with the ¹Department of Agriculture, Faculty of Applied Science, Cape Peninsula University of Technology, P.O. Box XB Wellington 7654, South Africa (e-mail: tangam@cput.ac.za)

A.T Adetunji is with the ¹Department of Agriculture, Faculty of Applied Science, Cape Peninsula University of Technology, P.O. Box XB Wellington 7654, South Africa (e-mail: adetunji@cput.ac.za)

F.B Lewu is with the ¹Department of Agriculture, Faculty of Applied Science, Cape Peninsula University of Technology, P.O. Box XB Wellington 7654, South Africa (e-mail: lewuf@cput.ac.za)

collection commenced four weeks after transplanting. The total yield was recorded at harvest for both seasons.

D. Statistical analysis

Analysis of variance ANOVA at 95% confidence limit and comparison of means was carried out on vegetative growth and yield. Analysis of data of the plant data collected using IBM SPSS software (2020). Means separation was done using Fisher's Least Significant Difference (LSD).

III. RESULTS AND DISCUSSION

A. Effect of Phosphorus Fertilizer Treatment on Vegetative Growth and Yield of *Brassica oleracea*.

TABLE I: INFLUENCE OF FERTILIZER TREATMENT ON VEGETATIVE GROWTH AND YIELD OF CABBAGE 2019

Treatment	Number of leaves	Plant height (cm)	Leaf length (cm)	Leaf width (cm)	Yield (kg/ha)
T1	13.13 ^a	20.62 ^a	15.73 ^a	13.38 ^a	1.53 ^a
T2	12.71 ^a	21.03 ^a	15.72 ^a	14.07 ^a	1.96 ^a
T3	12.42 ^a	21.13 ^a	13.29 ^a	12.35 ^a	2.01 ^a
T4	13.13 ^a	21.89 ^a	16.38 ^a	13.98 ^a	2.20 ^a
LSD	NS	NS	NS	NS	NS

Means in the same column with the same superscript are not significantly different ($P>0.05$). LSD= Least significant difference; NS= Not Significant; T1= P 0%, T2= P 50%, T3= 75%, T4= 100%.

Phosphorus (P) fertilizer treatment had no significant ($p>0.05$) effect on vegetative growth and yield as shown in (Table I & II). This could be due to that only P was varied in the study and the other two significant macronutrients nitrogen and potassium we kept constant. However, the application of phosphorus fertilizer at 222.5 kg/ha (T3) in the year 2019 increased plant height, leaf length, leaf width and yield by 6, 4, 4, and 44% compared to the control. These findings were in contrast with those of [16] who reported that phosphorus significantly increased vegetative growth, yield and also improved soil P content. Soil results from this study showed an increase in P content before the second season of planting which increased from 23 to 32mg/kg. The ideal soil pH that is suitable to cultivate cabbage ranges from 5.5 to 6.5 and initial soil analysis results for this field experiment was 4.6 and might have restricted phosphorus availability to plants. These observations are in line with those of [1], as the results showed that in acidic soils phosphorus is fixed by aluminium and iron.

TABLE II: INFLUENCE OF FERTILIZER TREATMENT ON VEGETATIVE GROWTH AND YIELD OF CABBAGE 2020

Treatment	Number of leaves	Plant height (cm)	Leaf length (cm)	Leaf width (cm)	Yield (kg/ha)
T1	16.03 ^a	17.01 ^a	15.79 ^a	15.62 ^a	5.02 ^a
T2	15.64 ^a	16.93 ^a	15.13 ^a	14.95 ^a	5.82 ^a
T3	16.45 ^a	18.89 ^a	17.58 ^a	16.84 ^a	5.86 ^a
T4	15.00 ^a	16.48 ^a	15.35 ^a	14.94 ^a	5.97 ^a
LSD	NS	NS	NS	NS	NS

Means in the same column with the same superscript are not significantly different ($P>0.05$). LSD= Least significant difference; NS= Not Significant; T1= P 0%, T2= P 50%, T3= 75%, T4= 100%

Similar results were obtained during the year 2020, phosphorus fertilizer treatment had no significant effect ($p>0.05$) on all measured vegetative parameters. An increase in yield during the second season harvest of cabbage was observed even though not significant. The highest yield of 5.97kg/ha was obtained during the second year of planting compared to 2.20kg/ha on the first year.

B. Effect of treatment on vegetative growth measured at different dates.

The different dates of data collection had a significant difference in the vegetative parameters measured for the two growing seasons (Table III & IV). This is due to the general growth pattern of plants that affects the vegetative parameters measured and increases as the season progresses. The date in which parameters were measured had a significant ($p<0.05$) effect on all vegetative parameters measured for both seasons. These findings are in agreement with those of [5], who reported an increase in plant height and number of leaves per plant at six week after transplanting. Reference [3], stated that different vegetables differ in their ability to uptake phosphorus which eventually affects the general growth of the crop.

TABLE III: INFLUENCE OF TIME ON DATA COLLECTED ON VEGETATIVE GROWTH DURING THE YEAR 2019.

Period	Number of leaves	Plant height (cm)	Leaf length (cm)	Leaf width (cm)
t1	8.75 ^c	14.68 ^b	8.26 ^c	7.48 ^c
t2	13.39 ^b	24.06 ^a	17.73 ^b	14.14 ^b
t3	16.41 ^a	24.83 ^a	19.85 ^a	18.71 ^a
LSD	3.02	9.38	2.12	4.57

Means in the same column with the same superscript are not significantly different ($P>0.05$). LSD= Least significant difference; t1= First date of data collection, t2= Second date of data collection, t3= Third date of data collection.

TABLE IV: INFLUENCE OF TIME ON DATA COLLECTED ON VEGETATIVE GROWTH DURING THE YEAR 2020.

Period	Number of leaves	Plant height (cm)	Leaf length (cm)	Leaf width (cm)
t1	11.28 ^c	11.89 ^c	10.71 ^b	9.67 ^b
t2	15.74 ^b	20.77 ^a	18.47 ^a	18.27 ^a
t3	20.32 ^a	19.33 ^b	18.71 ^a	18.83 ^a
LSD	4.46	1.44	0.24	0.65

Means in the same column with the same superscript are not significantly different ($P>0.05$). LSD= Least significant difference; t1= First date of data collection, t2= Second date of data collection, t3= Third date of data collection.

IV. CONCLUSION AND RECOMMENDATIONS

The availability of phosphorus (P) to plants can be restricted due to a number of factors such as crop type and soil condition such as the soil pH. The findings from the field experiment indicated that P did not affect growth parameters measured, however, an increase in yield was observed in the second year of the experiment. Research on macronutrients has mainly focused on nitrogen which influences vegetative growth and formation of stems and the findings from the study indicated that P had no significant effect on parameters measured. So further research on fertilizer combination is essential, including potassium and other macronutrients to provide the recommended fertilizer ratio that will improve production of cabbage in Western Cape, Wineland region.

ACKNOWLEDGMENT

The authors would like to thank the Cape Peninsula University of Technology (CPUT) for providing support and research facilities and special gratitude to the head of department in Agriculture in the institution for financial support for the duration of the research.

REFERENCES

- [1] Ch'ng, H.Y., Ahmed, O.H., Majid, N.M. 2014. Improving phosphorus availability in an acid soil using organic amendments produces from agroindustrial wastes. *The Scientific World Journal*, 1-6.
- [2] DAFF (Department of Agriculture Forestry and Fisheries). n.d. Production guidelines for cabbage. <https://www.nda.agric.za/docs/brochures/prodguidecabbage.pdf>
- [3] Fohse, D., Claassen, N., Jungk, A. 1988. Phosphorus efficiency of plants. *Plant and Soil*, 110: 101-109.
- [4] Halim, G.M.A., Ahmed, M.S., Islam, A., Hossain, S. 1994. Effect of different doses of NPK on growth and yield of cabbage. *Ann. Bangladesh Agriculture*, 4(2): 157-160.
- [5] Kartika, J. G., Herdyanti, S.T., Fadilah, N. 2017. Improving growth and production of cabbage (*Brassica oleraceae* L.) with compound fertilizer application. *Journal of Tropical Crop Science*, 4(2): 58- 63.
- [6] Khan, R., Ahmed, S., Khan, S., Ahmed, F., Zaman, M., Khan, B.A. 2002. Effect of different levels of nitrogen, phosphorus and potassium on the growth and yield of cabbage. *Asian Journal of Plant Sciences*, 1(5): 548-549.
- [7] Maseko, I., Mabhaudhi T., Tesfayi S., Avaya, H.T., Fezzehazion, M., Du Plooy, C.P. 2018. African leafy vegetables: A review of status, production and utilisation in South Africa. *Sustainability*, (10) 16, 1-16.
- [8] Muleke, A.M., Saidi, M., Itulya, F.M., Martin, T., Ngouajio, M. 2013. The assessment of the use of eco- friendly nets to ensure sustainable cabbage seedling production in Africa. *Agronomy*, 3: 1- 12.

- [9] Nikzad, M., Kumar, J.S.A., Anjanappa, M., Amarananjundeswara, H., Dhananjaya, B.N., Basavaraj, G. 2020. Effect of fertigation, levels on growth and yield of cabbage (*Brassica oleracea* L. var. capitata). *International Journal of Current Microbiology and Applied Sciences*, 9 (1): 1240- 1247.
- [10] Tanga, M., Lewu, F.B., Oyedeji, A.O., Oyedeji, O.O. 2020. Growth, yield and phytochemical constituents of *Arctium lapp* L. in response to phosphorus and potassium fertilizer application. *18th SOUTH AFRICA Int'l Conference on Agricultural, Chemical, Biological & Environmental Sciences* (ACBES-20) Nov. 16-17, 2020 Johannesburg (SA): 352-357.
- [11] Fatima, S.N., Norida, M., Zaharah, S.S. 2019. Effect of different nitrogen fertilization on cabbage (*Brassica oleracea*) and development of diamondback moth (*Plutella xylostella*). *Food Research*, 3(4): 342-347.
- [12] Kerr, B. 2014. Nitrogen: the key to high cabbage yield. *Farmers Weekly*. <https://www.farmersweekly.co.za>.
- [13] Yeshiwas, Y. 2017. Effect of different rate of nitrogen fertilizer on the growth and yield of cabbage (*Brassica oleraceae*) at Debre Markos, North West Ethiopia. *African Journal of Plant Science*, 11(7): 276-281.
- [14] Haque, K.M.F., Jahangir, A.A., Haque, M.E., Mondal, R.K., Jahan, M.A.A., Sarker, M.A.M. 2006. Yield and nutritional quality of cabbage as affected by nitrogen and phosphorus fertilization. *Bangladesh J. Sci. Ind. Res.*, 41(1-2): 41-46.
- [15] Atanasova, E., Mitova, I., Dimitrov, I., Stancheva, I. 2007. Effect of different fertilizer sources on the quality of head cabbage. *Journal of Applied Horticulture*, 9(1): 74-76.
- [16] Filho, A.B.C., Silverio da Silva, G., Cortez, J.W.M. 2013. Phosphorus fertilization of 'Fuyutoto' cabbage in phosphorus-rich Eutrustox soil. *Chilean Journal of Agricultural Research*, 73(3): 288-292.